

The effect of adding hip strengthening exercises to lumbar stabilizing exercises on gait for the treatment of non-specific low back pain: a randomized controlled trial

DOI: <https://doi.org/10.5114/pq/175678>

Amir Mohamed Saleh¹ , Nabil Mahmoud Abdel-Aal¹ , Ahmed Bayuomy Abo Elatta² 

¹ Department of Basic Sciences, Faculty of Physical Therapy, Cairo University, Giza, Egypt

² Department of Physical Therapy, Police Authority Hospital, Cairo, Egypt

Abstract

Introduction. To evaluate the impact of adding hip strength exercises to lumbar stabilizing exercises on pain, disability, and spatio-temporal parameters of gait in the treatment of non-specific low back pain (NSLBP).

Methods. In this randomized controlled trial, 60 patients diagnosed with NSLBP were randomized into two equal groups. The subjects were randomly allocated to either group A, the control group ($n = 30$), performing lumbar stabilizing exercises, or group B, the study group ($n = 30$), receiving lumbar stabilizing exercises and a progressive hip strengthening exercise program. The frequency of intervention was 3 sessions per week for six weeks. In this trial, the Visual Analogue Scale for pain intensity, Modified Oswestry Disability Index for disability, and Kinovea 2D motion analysis for spatio-temporal parameters were evaluated pre-treatment and after six weeks.

Results. There were statistically significant differences between groups in changes in pain scores ($p = 0.035$), the disability index ($p = 0.012$), and COG vertical displacement ($p = 0.017$). There were no statistically significant differences in stride length, stride time, stride speed, or cadence between the control and study groups ($p > 0.005$).

Conclusions. Adding hip strengthening exercises to a lumbar stabilization exercise program can reduce pain and disability and decrease COG vertical displacement more than a lumbar stabilization exercise program alone in NSLBP patients. However, no differences were found in gait speed, cadence, stride length, and stride time.

Key words: non-specific low back pain, Kinovea, hip strengthening exercises, and lumbar stabilization exercises

Introduction

As one of the most common musculoskeletal conditions, low back pain (LBP) affects more than 80% of the global population and is a leading cause of lost work time, doctor visits, and decreased quality of life [1]. Approximately 10% to 40% of patients with LBP progress to develop chronic LBP [2, 3]. Most cases of LBP fall under the category of non-specific LBP (NSLBP), which has no identifiable disease or underlying cause and affects 90% of the LBP population [4].

Weakness of the superficial trunk as well as abdominal muscles is an essential contributing factor to LBP. So, strengthening these muscles is usually linked with major reductions in LBP and lower functional impairment [5]. Patients with LBP have insufficient activation of the lumbar multifidus (LM), transversus abdominis (TrA), as well as obliquus internus (OI) muscles, all of which contribute to lumbar spinal stability [6].

NSLBP has a significant influence on proprioception [7] and the spatio-temporal parameters of gait [8, 9]. The findings can be used in selecting better rehabilitation procedures [8]. Walking patterns in elderly people with LBP are significantly distinct from those in older adults without LBP, even after controlling for age and gender. The wider the steps, the higher the risk of LBP when walking quickly [9].

Clinical practice guidelines suggest exercise therapy as an appropriate therapeutic option for those suffering from NSLBP. Patients with CLBP can benefit greatly from a program of segmental stabilization as well as strengthening exercises [10, 11]. Patients having LBP commonly suffer from a lack of

hip abductor strength [12]. Muscle strengthening exercise, which may involve lumbar exercises, is a great addition to exercise treatment programs for individuals with LBP for the purpose of rehabilitation as well as preservation of easy daily living [13]. Core muscle dysfunction in NSLBP can contribute to poor motor control during gait which may lead to compensatory movement [14].

Although lumbar stabilizing exercises and progressive hip strengthening exercise programs were already tested for their effect in managing LBP, up to the researchers' knowledge, there is a gap in the body of knowledge about their effectiveness on spatio-temporal gait parameters. So, the objective of the study was to evaluate the impact of progressive hip strengthening exercises when applied in combination with lumbar stabilizing exercises on pain, disability, and spatio-temporal gait parameters in patients with NSLBP.

Subjects and methods

This study was a randomized controlled trial. Patients were recruited from the physical therapy outpatient clinic of Cairo University. From December 2021 to December 2022, patients were evaluated in a physical therapy clinic where they received treatment.

Subjects

Eligibility was determined for all individuals with NSLBP who were referred for physiotherapy by a neurologist. Patients

Correspondence address: Ahmed Bayuomy Abo Elatta, Department of Physical Therapy, Police Authority Hospital, Taha Hussin Street, New Cairo, Cairo P.O. 4731501, Egypt, e-mail address: ahmedbayuomy@gmail.com; <https://orcid.org/0009-0009-2050-6061>

Received: 10.06.2023

Accepted: 22.11.2023

Citation: Saleh AM, Abdel-Aal NM, Elatta ABA. The effect of adding hip strengthening exercises to lumbar stabilizing exercises on gait for the treatment of non-specific low back pain: a randomized controlled trial. *Physiother Quart.* 2025;33(1):42–49; <https://doi.org/10.5114/pq/175678>.

were included if they experienced pain below the level of T12 but not below the buttock line, were between the ages of 18 and 60, and had chronic nonspecific LBP (greater than 12 weeks). Cancer patients, those with spinal cord injuries or spinal osteoporosis, those with primary joint diseases including active rheumatoid arthritis (RA) or metabolic bone disorder, and those with significant psychological problems were not included.

Sample size

To reduce the probability of type II errors, the sample size was calculated before the study began. Using an effect size of 0.41, a power analysis of 80%, and a two-tailed significance level of 5%, G*Power (version 3.1.9.2; Germany) determined the necessary sample size for the study. It was originally expected that there would be 49 patients with NSLBP, but after considering dropouts, the overall sample size was calculated to be 60.

Randomization

Sixty people with NSLBP were randomized between two groups (A and B). A random block generation computer tool, found at <http://www.randomization.com/>, was used to assign groups at random. To reduce the potential for bias and variability among the two groups, patients were randomly assigned to blocks of four, six, and eight with an allocation ratio of 1:1. The author, who had no role in participant selection, data collection, or treatment, used opaque, sealed envelopes to make the concealed allocation. Another author applied baseline and 6-week measurements. Finally, the envelopes were opened by an author who continued with the treatment consistent with group allocation.

Outcome measures

All outcome measures were evaluated at baseline and 6 weeks later for each treatment intervention.

– Pain intensity level: The pain intensity was measured utilizing a Visual Analog Scale (VAS). This measurement tool uses a transverse continuous line of 100 mm in length, beginning with no pain at all on the left side while the right side represents the worst pain imaginable. The patient is requested to express their degree of pain according to the scale. The VAS is generally regarded as a valid and reliable tool for pain measurement [15, 16].

– Disability assessment: The Arabic version of the modified Oswestry Disability Questionnaire (MODQ) was used. This index is a well-validated, self-reported questionnaire created for LBP with ten components in the Arabic version [17]. It is a valid and reliable tool for the assessment of patient function [18]. There are 10 questions on this self-evaluation questionnaire, and each one can be graded from 0 to 5. These include pain, self-care, lifting, moving, walking, sitting, standing, and sexual and social life, as well as travel and sleeping difficulties due to LBP. The total score is divided by 50 and then multiplied by 100, giving a resulting percentage of disability ranging from 0% (no disability) to 100% (total disability). Scores on this scale can be interpreted in the following ways: Below 20% impairment is considered minimal disability, between 20% and 40% is considered moderate disability, between 40% and 60% is considered severe disability, between 60% and 80% is considered debilitating LBP, and above 80% is considered excessive incapacity (bed-ridden) [19].

– Spatio-temporal assessment (Kinovea 2D free software, version 0.9.5): The spatio-temporal gait parameters include stride length, stride time, stride speed, cadence, and COG vertical displacement. Kinovea two-dimensional (2D) analyses have been shown to be a valid, reliable, accessible, low-cost, and free alternative to assess lower limb kinematics during gait [20, 21]. The goals and methods of the study were thoroughly discussed with each participant, and all instructions were given verbally. Reflective markers were applied to the anterior superior iliac spine, greater trochanter, lateral condyle of the femur, and lateral malleolus of the fibula [22].

Despite the fact that the system (Sony Camera-Kinovea) utilized is technically a markerless motion capture system, the markers help Kinovea focus on the action when it's time to track the action (play back the video) [23]. Each participant was instructed to walk at a normal pace along the 5-meter walking route. A high-speed digital video camera (Sony HXRNX5U NXCAM; Sony Corp., Minato, Tokyo, Japan) was used to record the gait. The camera was placed on a level tripod approximately 0.3048 m (1 ft) above the floor and at a distance of 2.43 m (8 ft) perpendicular to the center of the pathway [22, 23].

The walking protocol was performed 5 times, and each time a video was taken using a Sony camera to capture the action (with an emphasis on the walking gait). Marker placement and the remainder of the experiment took approximately 20 min in total for one individual [22]. Each recorded video was first transferred to a computer, where it was then played again to look for any evident problems. Furthermore, poor-quality captures were eliminated. For the accepted videos, Kinovea was used to find and identify all four major markers and to follow the movement of the markers during the whole clip (walking gait) [23].

To account for initiation and fatigue, the first and last cycles of gait were excluded from the study. Thus, only the intermediate (2–4) cycles were examined. Following the completion of the video tracking process, the software's capabilities were used to measure and record the gait spatio-temporal characteristics (speed, cadence, stride time, stride length, step length, step time, and COG vertical displacement) [20–23].

Intervention

Patients in each of the two groups performed the same set of exercises under the close supervision of a physical therapist. Each patient was scheduled for 3 sessions per week over a 6-week period. Each treatment visit lasted 30–45 min. All participants received an education session at the beginning.

Group A: control group

Patients received only lumbar stabilizing exercises. Each exercise required an isometric hold that lasted for 7–8 s. Ten repetitions of each exercise were performed, with 3 s of rest between sets. The patients rested for a full minute between each exercise [24, 25]. The description of lumbar stabilizing exercises is shown in Supplementary 1.

Group B: study group

Hip strengthening exercises

Patients received lumbar stabilization exercises in addition to progressive hip strengthening exercises. Progressive hip strengthening was done according to the Delorme protocol. First, the 10-repetition maximum was determined for

each patient. Then the therapist uses 50% of the 10-repetition maximum as the training load for the first 2 weeks, then 75% of the 10-repetition maximum in the second 2 weeks, then 100% of the 10-repetition maximum in the final 2 weeks. All exercises were performed 10 times per set for 2 sets per session. Three sessions per week for 6 weeks were prescribed. Each exercise was performed for 30 s at full isometric contraction, followed by 10 s of rest in the starting position [26]. The description of hip strengthening exercises is shown in Supplementary 2.

Statistical analysis

Shapiro–Wilk was used to evaluate for data normality. Levene’s test for homogeneity of variances was performed to examine the homogeneity among groups. The data followed a normal distribution, and the variances were the same. The subjects’ characteristics were compared between groups using an independent *t*-test. Analyses of treatment effects on VAS, MODI, COG vertical displacement, stride length, stride time, stride speed, and cadence were performed using a mixed MANOVA. For further multiple comparisons, post-hoc tests with the Bonferroni correction were performed. All statistical tests were performed at the $p < 0.05$ level of significance. The Windows version of the Statistical Package for the Social Sciences (SPSS) version 25 was used for the analyses.

Results

Baseline data as well as subject characteristics for both the control and study groups are presented in Table 1. Age, weight, height, and body mass index did not differ significantly ($p > 0.05$) between groups. In addition, baseline data did not vary significantly ($p > 0.05$) between the groups. The flow-chart of the study was presented in Figure 1.

Table 1. Comparison of age, weight, height BMI, and occupation between control and study groups

Characteristics	Control group (mean ± SD)	Study group (mean ± SD)	MD	<i>p</i> -value
Age (years)	37.56 ± 7.52	35.8 ± 6.86	1.76	0.34
Weight (kg)	80.3 ± 10.47	80.36 ± 10.76	-0.06	0.98
Height (cm)	175.5 ± 7.04	173.3 ± 7.22	2.2	0.23
BMI (kg/m ²)	26.04 ± 2.86	26.81 ± 3.78	-0.77	0.37
Occupation, <i>n</i> (%)				
teachers	9 (30%)	7 (23.33%)	$\chi^2 = 1.12$	0.77
office workers	6 (20%)	4 (13.33%)		
factory workers	8 (26.67%)	10 (33.34%)		
students	7 (23.33%)	9 (30%)		

A significant interaction between treatment and time was found using mixed MANOVA ($F = 8.97, p = 0.001, \eta^2 = 0.54$). The time factor was found to have a significant main effect ($F = 76.77, p 0.001, = 0.97$). The treatment main effect was statistically significant ($F = 2.82, p 0.01, \eta^2 = 0.27$). According to Table 2, there were no pre-treatment statistically significant differences among the groups. There was a significant decline in VAS, MODI, and COG vertical displacement in the study group when compared to that of the control group post-treatment ($p < 0.05$). No significant differences were detected in stride length, stride time, stride speed, and cadence between control and study groups post-treatment ($p > 0.05$, Table 3).




Figure 1. CONSORT flow chart

Table 2. Baseline clinical characteristics of the subjects (n = 60)

Variable	Control group (mean ± SD)	Study group (mean ± SD)	MD (95% CI)	p-value
VAS	54 ± 20.1	50 ± 17.61	4 (-5.76: 13.76)	0.416
MODI (%)	29.86 ± 9.51	29.7 ± 8.85	0.16 (-4.58: 4.91)	0.944
COG vertical displacement (cm)	5.88 ± 1.23	5.98 ± 1.35	-0.1 (-0.76: 0.57)	0.771
Stride length (cm)	136.13 ± 12.71	134.22 ± 10.56	1.91 (-4.12: 7.95)	0.528
Stride time (ms)	1294.94 ± 125.74	1256.63 ± 172.39	38.3 (-39.67: 116.29)	0.330
Stride speed (m/s)	1.11 ± 0.18	1.03 ± 0.24	0.08 (-0.04: 0.18)	0.206
Cadence (step/min)	92.14 ± 9.41	91.27 ± 11.51	-0.87 (-4.57: 6.29)	0.751

VAS – Visual Analogue Scale, MODI – Oswestry disability index, COG – centre of gravity, MD – mean difference
p-value < 0.05 indicates statistical significance

Table 3. Between-group comparisons after 6 weeks of interventions (n = 60)

Variable	Control group (mean ± SD)	Study group (mean ± SD)	MD (95% CI)	p-value	η ²
VAS	29.33 ± 12.36	23 ± 10.22	6.33 (0.46:12.19)	0.035	0.075
MODI (%)	21.8 ± 9.25	16.03 ± 7.97	5.77 (1.3:10.23)	0.012	0.103
COG vertical displacement (cm)	5.45 ± 1.09	4.77 ± 1.05	0.68 (0.12: 1.23)	0.017	0.094
Stride length (cm)	145.8 ± 12.69	148.57 ± 12.52	-2.77 (-9.28: 3.74)	0.398	0.012
Stride time (ms)	1246.26 ± 146.25	1195.16 ± 141.56	51.1 (-23.28: 125.48)	0.174	0.032
Stride speed (m/s)	1.19 ± 0.21	1.25 ± 0.15	-0.06 (-0.16: 0.02)	0.153	0.035
Cadence (step/min)	97.61 ± 11.77	99.97 ± 12.21	-2.36 (-8.56: 3.83)	0.449	0.010

VAS – Visual Analogue Scale, MODI – Oswestry disability index, COG – centre of gravity, MD – mean difference
p-value < 0.05 indicate statistical significance

Table 4. Within groups changes pre-post intervention of control and study groups

Outcome	Control group		Study group	
	change from baseline to 6 weeks		change from baseline to 6 weeks	
	MD (95% CI)	p-value	MD (95% CI)	p-value
VAS	24.67 (19.56: 29.77)	0.001	27 (21.89: 32.1)	0.001
MODI (%)	8.06 (5.85, 10.27)	0.001	13.67 (11.45: 15.87)	0.001
COG vertical displacement (cm)	0.43 (0.11: 0.73)	0.008	1.21 (0.89: 1.51)	0.001
Stride length (cm)	-9.67 (-12.43: -6.89)	0.001	-14.35 (-17.12: -11.58)	0.001
Stride time (ms)	48.68 (5.59: 91.76)	0.02	61.47 (18.38: 104.55)	0.006
Stride speed (m/s)	-0.08 (-0.14: -0.02)	0.009	-0.22 (-0.28: -0.16)	0.001
Cadence (step/min)	-5.47 (-8.55: -2.39)	0.001	-8.7 (-11.78: -5.62)	0.001

VAS – Visual Analogue Scale, MODI – Oswestry disability index, COG – centre of gravity
p-value < 0.05 indicates statistical significance

There was a significant decline in VAS, MODI, COG vertical displacement, and stride time post-treatment compared to pre-treatment in the control and study groups (p > 0.05). There was a significant increase in stride length, stride speed, and cadence post-treatment compared to pre-treatment in the control and study groups (p > 0.01) as shown in Table 4.

Discussion

The present study aimed to examine the impact of adding hip strengthening exercises to the lumbar stabilization pro-

gram on pain, disability, and the spatio-temporal parameters of gait in NSLBP patients. The current study's results found significant improvement in pain, functional level, and spatio-temporal parameters of gait in both groups for all variables (p < 0.05). A significant difference has been detected between the control group and the study group in pain, disability, and COG vertical displacement favouring the study group (p < 0.05). However, there were no statistically significant differences between groups for stride length, stride time, stride speed, and cadence in patients with NSLBP (p > 0.05).

The findings of the present study could be explained by many reasons. First, hip muscle weakness and hip muscle

imbalance lead to lumbar pain in NSLBP. It has also been suggested that weak hip extensors and abductors contribute to LBP [27]. In addition, piriformis tightness from a lack of hip extensor strength contributes to LBP and limited hip rotatory motion [28]. As a result, it has been highlighted how essential it is to strengthen the hip extensors as well as abductors to reduce LBP [29]. Also, strengthening exercise can lead to improvement in functional scores which has an effect on pain as it disturbs a vicious cycle of pain sensitization and reinforcing disability [30]. Hip strengthening has been shown to improve the range of motion, decrease pain, enhance strength, and make movement easier for those with limited mobility [30]. Confidence improved and depression and fear related to LBP diminished when moving around without pain [31]. Therefore, patients can easily perform ADLs when they can walk around without experiencing pain [31]. Improvements in spinal stability as well as decreased stress on the spine were seen in all participants when progressive hip strengthening exercises were added to lumbar stabilization exercises. This helped reduce the impairment encountered during everyday activities [32]. Muscle imbalance as well as abnormal movement patterns developed as a protective measure for injured areas, but they only increased pain in their daily activities [31, 32].

Many studies [28, 30, 33, 34] reported that adding hip strengthening exercises to the lumbar stabilization program improved disability and pain in patients with NSLBP. Similar to the findings of the current study, Lee et al. [33] found a significant decline in pain in patients having LBP following 6 weeks of lumbar stabilization exercises.

However, this study's findings contradicted those of previous studies [35, 36]. Kendall et al. [35] conducted a study on NSLBP patients and found no significant improvement in pain and disability. The reason for this contradiction may be due to a reduced treatment session frequency as all participants received only one session per week. Fukuda et al. [36] concluded there were no significant differences in adding hip strength exercise to pain intensity and functional disability in NSLBP. The difference in the nature, frequency, and duration of treatment could account for this disparity.

In the current study, hip strengthening exercises reduced the COG significantly. The mechanism by which hip strengthening exercise induced that effect could be explained by the fact that the thoracolumbar fascia connects the hip muscles to the contralateral latissimus dorsi in the lower back, allowing the lumbar spine to act as a lever for transferring energy as well as loads to the lower limbs.³⁷ Furthermore, the gluteal muscle group maintains pelvic stability, which in turn supports the spine. Therefore, individuals with LBP may benefit greatly from doing hip strengthening exercises in addition to their regular strength training for the trunk musculature [37, 38].

The gluteus medius makes up around 60% of the abductor's cross-sectional area, making it the biggest of the abductor muscles. Because the gluteus medius plays an essential part throughout the single-leg stance time period, strengthening the hip abductors through targeted strength training is essential for performing functional lower limb motions. The lateral pelvis' dynamic stability is largely controlled by the gluteus medius muscle [39], thus facilitating normal movement patterns, thereby improving balance by decreasing the vertical COG displacement [29, 35]. When the center of gravity (COG) is greatly altered, such as in a single-leg stance, the hip muscles, especially the gluteus maximus, are responsible for preserving pelvic stability as well as regulating the rotational motion of the lower extremities. Therefore, abnormal segmental motion of the lumbar spine during walking or standing

may result from a lack of strength in these muscles, resulting in poor pelvic stability [38].

The findings of the present study were confirmed by many studies [29, 40, 41] demonstrating that the addition of hip strengthening exercises to the lumbar stabilization program improved normal movement patterns, dynamic stability, and decreased vertical COG displacement in patients with NSLBP. Similarly, a study carried out by Do and Yim [42] found that strengthening exercises for the hips not only improved single-stance performance, but also helped restore normal gait patterns and improved the strength of weak hip muscles, all of which had a positive effect on lumbar stability.

On the contrary, the study findings come in contradiction with other studies [36, 43]. Fukuda et al. [36] did not support the positive effect of hip strengthening exercises on either clinical or kinematic outcomes in patients with NSLBP. This contradiction may be attributed to the difference in kinematic measuring tools and the nature, frequency, and duration of intervention. Bennell et al. [43] demonstrated a decline in pain and improvement in function after hip strengthening but detected no changes in mechanics during walking and joint loading while walking. This disparity is due to different methods of assessment and different populations.

Limitations

This study had limitations. First, the number of patients in this study was relatively limited which might pose a risk to positive findings. Second, the study participants were male subjects only. Additional studies are recommended to amend these limitations. Third, it would be helpful to follow the long-term impacts of this intervention, but we lack sufficient data to do so. Additional studies should aim to address this. Furthermore, the back muscle activities have not been investigated in the current study. Therefore, further studies should investigate muscle activity changes using electromyography.

Conclusions

Adding hip strengthening exercises to lumbar stabilization exercise programs can reduce pain, disability, and decrease COG vertical displacement more than lumbar stabilization exercise programs alone in NSLBP patients. However, no differences were found in gait speed, cadence, stride length, or stride time.

Recommendations

From this trial, it is recommended that further studies compare males and females and undertake the use of larger sample sizes. For future research, it is proposed that a trial be performed to investigate the long-term effect of hip strengthening in NSLBP patients. Further studies should investigate muscle activity changes using electromyography.

Ethical approval

The research related to human use has complied with all the relevant national regulations and institutional policies, has followed the tenets of the Declaration of Helsinki, and has been approved by the Physical Therapy Research Ethics Committee (approval No.: P.T. REC/012/002997). The study's registration number is PACTR202112720946276 with the Pan African Clinical Trials Registry.

Informed consent

Informed consent has been obtained from all individuals included in this study.

Disclosure statement

No author has any financial interest or received any financial benefit from this research.

Conflicts of interest

The authors state no conflicts of interest.

Funding

This research received no external funding.

References

- [1] Hoy D, Brooks P, Blyth F, Buchbinder R. The epidemiology of low back pain. *Best Pract Res Clin Rheumatol*. 2010;24(6):769–81; doi: 10.1016/j.berh.2010.10.002.
- [2] Lee SW, Kim SY. Effects of hip exercises for chronic low-back pain patients with lumbar instability. *J Phys Ther Sci*. 2015;27(2):34–48; doi: 10.1589/jpts.27.345.
- [3] Mesner SA, Foster NE, French SD. Implementation interventions to improve the management of non-specific low back pain: a systematic review. *BMC Musculoskelet Disord*. 2016;17:258; doi: 10.1186/s12891-016-1110-z.
- [4] Hartvigsen J, Hancock MJ, Kongsted A, Louw Q, Ferreira ML, Genevay S, Hoy D, Karpainen J, Pransky G, Sieper J, Smeets RJ, Underwood M; Lancet Low Back Pain Series Working Group. What low back pain is and why we need to pay attention. *Lancet*. 2018;391(10137):2356–67; doi: 10.1016/S0140-6736(18)30480-X.
- [5] Ferreira PH, Ferreira ML, Maher CG, Refshauge K, Herbert RD, Hodges PW. Changes in recruitment of transversus abdominis correlate with disability in people with chronic low back pain. *Br J Sports Med*. 2010;44(16):1166–72; doi: 10.1136/bjism.2009.061515.
- [6] Butcher SJ, Craven BR, Chilibeck PD, Spink KS, Grona SL, Sprigings EJ. The effect of trunk stability training on vertical takeoff velocity. *J Orthop Sports Phys Ther*. 2007;37(5):223–31; doi: 10.2519/jospt.2007.2331.
- [7] Alasar S, Abdallah G, Elkabalawy M. Effect of proprioceptive training on lower back muscle performance in patients with chronic non-specific low back pain: a randomized controlled trial. *Physioth Q*. 2022;30(4):96–103; doi:10.5114/pq.2022.121155.
- [8] Najafi S, Rezasoltani Z, Abedi M. Effects of mechanical low back pain in spatiotemporal parameters of gait. *J Arch Milit Med*. 2019;6(4):e82816; doi: 10.5812/jamm.82816.
- [9] Hicks GE, Sions JM, Coyle PC, Pohlig RT. Altered spatiotemporal characteristics of gait in older adults with chronic low back pain. *Gait Posture*. 2017;55:172–6; doi: 10.1016/j.gaitpost.2017.04.027.
- [10] Wells C, Kolt GS, Bialocerkowski A. Defining Pilates exercise: a systematic review. *Complement Ther Med*. 2012;20(4):253–62; doi: 10.1016/j.ctim.2012.02.005.
- [11] França FR, Burke TN, Hanada ES, Marques AP. Segmental stabilization and muscular strengthening in chronic low back pain – a comparative study. *Clinics*. 2010;65(10):1013–7; doi: 10.1590/s1807-59322010001000015.
- [12] Cooper NA, Scavo KM, Strickland KJ, Tipayamongkol N, Nicholson JD, Bewyer DC, Sluka KA. Prevalence of gluteus medius weakness in people with chronic low back pain compared to healthy controls. *Eur Spine J*. 2016;25(4):1258–65; doi: 10.1007/s00586-015-4027-6.
- [13] Reiman MP, Weisbach PC, Glynn PE. The hips influence on low back pain: a distal link to a proximal problem. *J Sport Rehabil*. 2009;18(1):24–32; doi: 10.1123/jsr.18.1.24.
- [14] Van der Hulst M. Lumbar and abdominal muscle activity during walking in subjects with chronic low back pain: support of the hypothesis? *J Electromyogr Kinesiol*. 2010;20(1):31–8; doi: 10.1016/j.jelekin.2009.03.009.
- [15] Boonstra AM, Preuper HR, Reneman MF, Posthumus JB, Stewart RE. Reliability and validity of visual analogue scale for disability in patients with chronic musculoskeletal pain. *Int J Rehabil Res*. 2008;31(2):165–9; doi: 10.1097/MRR.0b013e3282fc0f93.
- [16] Alghadir AH, Anwer S, Iqbal A, Iqbal ZA. Test-retest reliability, validity, and minimum detectable change of visual analog, numerical rating, and verbal rating scales for measurement of osteoarthritic knee pain. *J Pain Res*. 2018;11:851–6; doi: 10.2147/JPR.S158847.
- [17] Fairbank JC, Pynsent PB. The Oswestry Disability Index. *Spine*. 2000;25(22):2940–52; doi: 10.1097/00007632-200011150-00017.
- [18] Algarni A, Ghorbels, Jones J, Guermazi M. Validation of an Arabic version of the Oswestry index in Saudi Arabia. *Ann Phys Rehabil Med*. 2014;57(9–10):653–63; doi: 10.1016/j.rehab.2014.06.006.
- [19] Fritz JM, Irrgang JJ. A comparison of a modified Oswestry Low Back Pain Disability Questionnaire and the Quebec Back Pain Disability Scale. *Phys Ther*. 2001;81(2):776–88; doi: 10.1093/ptj/81.2.776.
- [20] Damsted C, Larsen LH, Nielsen RO. Reliability of video-based identification of foot strike pattern and video time frame at initial contact in recreational runners. *Gait Posture*. 2015;42(1):32–5; doi:10.1016/j.gaitpost.2015.01.029.
- [21] Grigg J, Haakonssen E, Rathbone E, Orr R, Keogh JWL. The validity and intra-tester reliability of markerless motion capture to analyse kinematics of the BMX Supercross gate start. *Sports Biomech*. 2018;17(3):383–401; doi: 10.1080/14763141.2017.1353129.
- [22] Mathew J, Vanlalpeki T, Nair GG. Gait evaluation of institutionalized elders – a feasibility study. *Indian J Gerontol*. 2017;31(2):71–83.
- [23] Ya'akob Y, Hisham NAH, Patar MNA, Le CH, Mahmud J, Lee H, Yamamoto S, Herawati L. Walking Gait Analysis. Kinovea versus Motion Capture System. IEEE 13th Control and System Graduate Research Colloquium (ICSGRC), Shah Alam, Malaysia; 2022, pp. 187–91; doi: 10.1109/ICSGRC55096.2022.9845136.
- [24] Luque-Suárez A, Díaz-Mohedo E, Medina-Porqueres I, Ponce-García T. Stabilization exercise for the management of low back pain. *IntechOpen*. 2012;51(2):953–78; doi: 10.5772/34602.
- [25] McGill SM. *Low Back Disorders. Evidence-Based Prevention and Rehabilitation*. 3rd ed. Human Kinetics Publishers; 2015.
- [26] De Souza TP, Fleck SJ, Simao R, Dubas JP, Pereira B, De Brito Pacheco, da Silva AC, de Oliveira PR. Comparison between constant and decreasing rest intervals: influence on maximal strength and hypertrophy. *J Strength Cond Res*. 2010;24(7):1843–50; doi: 10.1519/JSC.0b013e3181ddae4a.
- [27] Beomryong K, Jongeun Y. Core stability and hip exercises improve physical function and activity in patients with non-specific low back pain: a randomized controlled trial. *Tohoku J Exp Med*. 2020;25(1):193–206; doi: 10.1620/tjem.251.193.
- [28] Yang S-J, Park K-N, Kyung M-S, Kim S-H. Comparison of hip rotation range of motion in jiu-jitsu athletes with and without low back pain. *Phys Ther Korea*. 2018;25(1):47–52; doi: 10.12674/ptk.2018.25.1.047.
- [29] Bade M, Cobo-Estevéz M, Neeley D, Pandya J, Gunderson T, Cook C. Effects of manual therapy and exercise targeting the hips in patients with low-back pain: a ran-

- domized controlled trial. *J Eval Clin Pract.* 2017;23(4): 734–40; doi: 10.1111/jep.12705.
- [30] Kang TW, Kim BR. The effects of stretching and strengthening exercise on the pain, pelvic tilt, functional disability index, and balance ability of patients with chronic lower back pain. *J Kor Phys Ther.* 2019;31(1):7–12; doi: 10.18857/jkpt.2019.31.1.7.
- [31] Telci EA, Yagci N, Can T, Cavlak U. The impact of chronic low back pain on physical performance, fear avoidance beliefs, and depressive symptoms: a comparative study on Turkish elderly population. *Pak J Med Sci.* 2013;29(2): 560–4; doi: 10.12669/pjms.292.3196.
- [32] Çelenay ST, Kaya DO. Effects of spinal stabilization exercises in women with benign joint hypermobility syndrome: a randomized controlled trial. *Rheumatol Int.* 2017;37(9):1461–8; doi: 10.1007/s00296-017-3713-6.
- [33] Lee HS, Kim DJ, Oh Y, Min K, Ryu JS. The effect of individualized gradable stabilization exercises in patients with chronic low back pain: case control study. *J Back Musculoskelet Rehabil.* 2016;29(3):603–10; doi: 10.3233/BMR-160724.
- [34] De Jesus FLA, Fukuda TY, Souza C, Guimarães J, Aquino L, Carvalho G, Powers C, Gomes-Neto M. Addition of specific hip strengthening exercises to conventional rehabilitation therapy for low back pain: a systematic review and meta-analysis. *Clin Rehabil.* 2020;34(11): 1368–77; doi: 10.1177/0269215520941914.
- [35] Kendall KD, Emery CA, Wiley JP, Ferber R. The effect of the addition of hip strengthening exercises to a lumbopelvic exercise program for the treatment of non-specific low back pain: a randomized controlled trial. *J Sci Med Sport.* 2015;18(6):626–31; doi: 10.1016/j.jsams.2014.11.006.
- [36] Fukuda TY, Aquino LM, Pereira P, Ayres I, Feio AF, de Jesus FLA, Neto MG. Does adding hip strengthening exercises to manual therapy and segmental stabilization improve outcomes in patients with nonspecific low back pain? A randomized controlled trial. *Braz J Phys Ther.* 2021;25(6):900–7; doi: 10.1016/j.bjpt.2021.10.005.
- [37] Hoffman SL, Johnson MB, Zou D, Van Dillen LR. Sex differences in lumbopelvic movement patterns during hip medial rotation in people with chronic low back pain. *Arch Phys Med Rehabil.* 2011;92(7):1053–9; doi: 10.1016/j.apmr.2011.02.015.
- [38] De Sousa CS, De Jesus FL, Machado MB, Ferreira G, Ayres IGT, de Aquino LM, Fukuda TY, Gomes-Neto M. Lower limb muscle strength in patients with low back pain: a systematic review and meta-analysis. *J Musculoskelet Neuronal Interact.* 2019;(19):69–78.
- [39] Baik S-M, Cynn H-S, Kim S-H. Understanding and exercise of gluteus medius weakness: a systematic review. *Phys Ther Korea.* 2021;28(1):27–35; doi: 10.12674/ptk.2021.28.1.27.
- [40] Baldrón RDM, Lobato DF, Carvalho LP, Wun PY, Santiago PR, Serrao FV. Effect of functional stabilization training on lower limb biomechanics in women. *Med Sci Sports Exerc.* 2012;44(1):135–45; doi: 10.1249/MSS.0b013e31822a51bb.
- [41] Jeong U-C, Sim J-H, Kim C-Y, Hwang-Bo G, Nam C-W. The effects of gluteus muscle strengthening exercise and lumbar stabilization exercise on lumbar muscle strength and balance in chronic low back pain patients. *J Phys Ther Sci.* 2015;27(12):3813–6; doi: 10.1589/jpts.27.3813.
- [42] Do, Yim. Effects of muscle strengthening around the hip on pain, physical function, and gait in elderly patients with total knee arthroplasty: a randomized controlled trial. *Healthcare.* 2020;8(4):489; doi: 10.3390/healthcare8040489.
- [43] Bennell KL, Hunt MA, Wrigley TV, Hunter DJ, McManus FJ, Hodges PW, Hinman RS. Hip strengthening reduces symptoms but not knee load in people with medial knee osteoarthritis and varus malalignment: a randomised controlled trial. *Osteoarthritis Cartilage.* 2010;18(5):621–8; doi: 10.1016/j.joca.2010.01.010.

Supplementary 1
 Abdominal stabilizing exercises

Exercise type	Description
Abdominal hollowing	The individual is supine, with his knees bent at a right angle. The pelvis and lower back are in proper alignment. The next step is for him to firmly contract his abdominal muscles in either a seated or kneeling position to elevate his internal pressure.
Side bridge	Under the patient's shoulder, their left forearm is resting on the floor. The individual raises his body into a "plank" or "side bridging" position. The forearm as well as the foot maintains their position to support. The same process is carried out on the right side of the body.
Supine extension bridge	The patient places both feet under their knees in a crock-lying position. Carefully supporting their weight using their shoulders instead of their necks, patients gradually raise their hips till their knees and shoulders are positioned in a straight line. After remaining in this position, the patient drops his hips to the floor gradually.
Straight leg rise from prone	The patient lies prone, supporting his head with his arms. Afterwards, he contracts the muscles in his right glute as well as hamstring and elevates his leg as much as he can toward the ceiling. Following remaining in this position for a while, the patient drops his leg gradually. The left side of the body receives a similar exercise.
Alternate arm and leg raise from quadruped	A quadruped position is performed by the individual. A cushion can be positioned under the knees for support if needed. The individual contracts his abdominal muscles for stabilization of his spine. The participant raises one arm as well as the opposite leg while keeping the abdominal muscles tense. After a few seconds, the participant lowers their arm and leg gradually and performs the exercise on the other side.
Prone bridge	While prone the elbow position is assumed by the patient. Then the patient pushes themselves up on their forearms, which are positioned between their shoulders and feet. The patient's hips and back remain in a straight, parallel position.

Supplementary 2
Progressive strengthening exercises

Exercise type	Description
Abductor strength	The patient is prone, with his or her hips and knees flexed to around 45 degrees. The therapist will stabilize the patient’s pelvis with their hands and forearms while the patient’s upper leg is extended and their ankle is neutral. The patient will then be instructed to lift their leg as much as possible while wearing a weighted cuff and to relax. Repeat the process with the other leg.
Extensor strength	The patient lies in a half-prone position, having the right leg firmly planted on the ground for support. After stabilizing the patient’s pelvis, the patient will be requested to elevate their left leg as much as they can while carrying a weighted cuff. Repeat the process with the other leg.
Flexors strength	The patient is lying supine, with his or her left leg extended. The next step is to raise the knee till the thigh is parallel to the ground. After stabilizing the patient’s pelvis, the patient will be requested to elevate their left leg as much as they can while carrying a weighted cuff. Repeat the process with the other leg.
Adductors strength	The patient should lie on their left side with their upper leg propped up by the therapist’s hand while the therapist’s opposite hand supports the patient’s pelvis. Have the patient raise their lower leg to their upper one while wearing a weighted cuff and hold the position. Repeat the process with the other leg.