

Therapeutic intervention on stabilisation parameters in patients with trigger points in suboccipital muscles: a case-control study

DOI: <https://doi.org/10.5114/pq/192350>

Miloslav Gajdoš¹, Wioletta Mikul'áková¹, Tomáš Uher¹, Čuj Jakub¹, Pavol Nechvatal¹, Lucia D. Kendrová¹

Department of Physiotherapy, Faculty of Health Care, University of Prešov, Presov, Slovak Republic

Abstract

Introduction. Hypertonus and the presence of trigger points (TrPs) in the suboccipital muscles (sOMs) frequently contribute to cervical proprioception disorders. Impaired cervical proprioception inevitably affects postural stability, which necessitates therapeutic intervention in conjunction with cervical spine therapy. The aim of this study was to examine the influence of the myofascial release technique (MFR) on stability parameters in patients diagnosed with TrPs in the sOM region.

Methods. Twelve patients (eight females and four males, mean age = 37 ± 9.5 years) underwent postural stability assessment using a force platform (Sensor Medica, Rome, Italy) before and after sOM treatment, with a time interval of one hour. The study's objectivity was supported by a control group comprising healthy participants (seven females and five males, mean age = 25 ± 3 years).

Results. The results indicated significant changes in stability parameters immediately after applying the MFR technique, particularly in patients with TrPs in the sOMs.

Conclusions. We recommend that patients undergoing MFR treatment maintain a safe position for at least a few minutes to prevent falls and dizziness.

Key words: posture, stability, cervical spine, proprioception, trigger points

Introduction

The suboccipital muscles (sOMs) primarily function to provide dynamic stability to the upper cervical spine. They perform small pendulum-like movements in the craniocervical junction, ensuring a neutral head position and offering proprioceptive feedback to the central nervous system [1]. The abundance of proprioceptors within the sOMs underscores their role in proprioception [2].

The highly delicate proprioceptive system of the cervical spine, which plays a crucial role in body posture and balance control [3], communicates the position of the head relative to the trunk, coordinates with the vestibular and visual systems, and is vital for the maintenance of body posture and balance [4]. Overloading the sOMs often presents as reflex changes, such as the development of myofascial trigger points (TrPs). These TrPs are characterised as hyperirritable spots within a muscle, manifesting as tender nodules palpable within taut bands [5, 6]. Upon palpation, these points are painful and may radiate pain unilaterally deep into the occipital region, as well as into the temporal area, eyes (ocular mask), and forehead. While patients typically localise the pain to the base of the skull, it is not directly associated with headache pain from the splenius cervicis muscle but rather with tension-type headaches [7, 8]. Disruption of cervical proprioception is a common issue in patients with neck pain, leading to disturbances in cervical sensorimotor control [2]. These disturbances can result in long-term changes in muscle tissue, exacerbate pain through peripheral and central nervous system mechanisms, sensitise the system, and contribute to dysfunctional movement patterns [1]. Moreover, muscle pain affects motor control strategies through central mechanisms [9, 10]. Research has shown that structural and functional alterations in the sOMs can influence muscle spindle firing or sensitivity, affect-

ing afference and leading to the aforementioned proprioceptive changes. As a result, flawed proprioception distorts the direct linear interaction between cervical proprioception and vestibular information, leading to an imprecise, subjective body orientation and spatial psychological representation (somatognosia). This can manifest as dizziness or a subjective sense of instability [11].

Examination of trigger points in the suboccipital muscles

Individuals with TrPs in the sOMs often experience headaches and exhibit a forward head posture, which is a common abnormality observed in clinical conditions affecting the cervical spine. The extension of the upper cervical spine in this posture results in the compression of cranio-cervical structures, especially the suboccipital and posterior cervical muscles. Visual assessment is a common method for evaluating an individual's body posture, with numerous studies indicating that manual evaluation of upper cervical dysfunction can assist in the precise diagnosis of patients experiencing cervical headache pain [12–14]. Proper manual assessment is crucial for distinguishing between TrPs in the sOMs and dysfunction of the craniocervical joint. It is important to note that the area just below the skull, where the sOMs are located, is highly sensitive to palpation, and therapists should be mindful of the pressure applied. Since the sOMs muscles are overlapped by the upper trapezius and other posterior neck muscles, they cannot be directly palpated. However, applying pressure to the anatomical projection of the sOMs can elicit pain specific to these muscles, distinct from the pain experienced in the splenius capitis or upper trapezius muscles. Additionally, palpation of each sOM may evoke slightly different pain patterns in some patients. Fernández de las Peñas

Correspondence address: Miloslav Gajdoš, Department of Physiotherapy, Faculty of Health Care, University of Prešov, Partizánska 1, 080 01 Presov, Slovak Republic, e-mail: miloslav.gajdos@unipo.sk; <https://orcid.org/0000-0002-6457-7867>

Received: 22.03.2024

Accepted: 15.08.2024

Citation: Gajdoš M, Mikul'áková W, Uher T, Jakub Č, Nechvatal P, Kendrová LD. Therapeutic intervention on stabilisation parameters in patients with trigger points in suboccipital muscles: a case-control study. *Physiother Quart*. 2025;33(3):46–53; doi: <https://doi.org/10.5114/pq/192350>.

et al. [15] developed a protocol for diagnosing TrPs in the sOMs, which includes assessing sensitivity in the suboccipital area, pain elicited by sustained pressure, and increased pain radiation during muscle contraction with upper cervical spine extension.

Assessment of sensorimotor control

According to the existing evidence, it is advisable for individuals with neck pain to undergo evaluation and treatment for impairment in cervical proprioception and disorders in sensorimotor control [16]. There are eight tests available to evaluate neck sensorimotor control in patients with chronic neck pain, including the Joint Position Error (JPE) test, posturography assessment, Subjective Visual Vertical test, Head Tilt Response test, Fly test, Smooth Pursuit Neck Torsion test, Head Stiffness test, and Rod-and-Frame test [3]. These tests may include assessing various subsystems involved in sensorimotor control, such as the oculomotor and vestibular systems, in addition to measuring cervical sensorimotor control [17].

Posturography measurement is a time-efficient technique that has proven useful in detecting significant changes in body sway during challenging situations. The implementation of force spectrum analysis in posturography testing can be a valuable tool for improving the evaluation of cervical spine function and monitoring rehabilitation treatments [18]. Head movements are also utilised in posturography measurements, which, according to several authors, significantly increase the movement of the centre of pressure (CoP) of the reacting force on the support surface [19, 20].

Treatment of trigger points in suboccipital muscles, neck pain, and headache

Various treatment approaches are used for cervicogenic headaches associated with TrPs in the sOMs. These approaches include electrotherapy, kinesiotherapy, and manual techniques [21]. Manual techniques employed for trigger point inactivation include ischaemic compression, combined trigger point compression with active muscle contraction, spray and stretch, post-isometric relaxation, and neuromuscular approaches [22]. Evaluating the most effective manual approaches poses a challenge since most therapists adopt a multimodal approach [23]. Corrective measures involve postural correction, ergonomic assessment, strengthening exercises, and self-stretching. Karegeannes et al. [16] recommend maintaining neck warmth, adjusting head position during daily and work activities, and ensuring proper overall body posture. They suggest stretching short and long neck extensors with manual patient assistance and placing a hand under the occipital bone, although manual therapy is not specifically mentioned. Though manual therapy targeting TrPs in the sOMs and related headache pain might yield positive outcomes in terms of pain intensity and frequency, additional research is needed to enhance the evidence regarding the impact of manual therapy on individuals suffering from tension-type headaches [24]. Specifically, the myofascial release technique (MFR) focuses on releasing shortened muscles and tension [25]. MFR utilises gentle pressure and stretching through direct and indirect approaches with the intention of reducing pain, optimising muscle length, and releasing fascial restrictions caused by injury, stress, or repeated static loading [26]. According to Ramezani et al. [25], the application of MFR and regular exercises leads to a rapid and early reduction in pain intensity, frequency, and pressure pain threshold in patients with cervicogenic headaches. As mentioned earlier, treatment

of proprioceptive disorders and sensory-motor control disturbances should be considered in patients with neck pain as it can have a positive impact on improving proprioception in the cervical spine area, chronic neck pain, cervical spine and balance function, and cervicocranial joint function [27]. Proprioception in the cervical spine area may improve through pain modulation or reduction, although research has yet to confirm this [3]. However, it is important to reduce muscle tension in the neck caused by pain to improve postural functions, which may lead to the normalisation of proprioceptive signals from neck muscles and a reduction in sensory-motor mismatch. Specific local treatment for neck pain relief that improves sensory-motor function is not discussed [3].

Objective of the study

Our study investigated the immediate impact of MFR on postural stability in patients with TrPs in the sOMs. While the effect of MFR on neck pain is well-known, the influence of MFR on the current state of proprioception and postural stability remains uncertain, and this was the primary focus of our study.

Subjects and methods

Characteristics of the study sample

The study included 12 patients diagnosed with TrPs in the sOMs (eight females and four males aged 37 ± 9.5). Participants were selected from patients seeking rehabilitative treatment for headaches associated with suboccipital pain and identified TrPs in these muscles. We used power analysis to determine the sample size, ensuring a sufficient number of participants for statistical validity. To ensure that all patients met the inclusion criteria for the study, they underwent a cervical spine examination conducted by a single experienced physiotherapist with over 15 years of practice. This examination included a patient interview, detailed medical history taking, a review of the patient's entire medical documentation and findings, if available, and a palpation examination of the sOMs.

Based on the CARE (case report) guidelines, we structured and wrote this case report to ensure a systematic and transparent description of the clinical case. The inclusion criteria for patient enrolment were adults within a specified age range, a forward head posture, TrPs in the sOMs with referred pain elicited by sustained manual pressure, a sensation of instability while walking but not true vertigo, and headaches occurring at least once a week in the last month. These criteria ensured that the study focused on the target population with relevant clinical characteristics. Exclusion criteria for the experimental group of patients were the manifestation of autonomic system symptoms, vertigo, or visual disturbances, severe distinct neck pain attributed to a herniated disc, spinal canal stenosis, or cervical spondylosis, any condition that might contraindicate the use of the MFR technique in the upper cervical region, and undergoing physiotherapy for neck and headache pain within the past week.

The control group consisted of 12 healthy individuals (7 females and 5 males aged 25 ± 3) without any headaches, neck pain, musculoskeletal disorders of the cervical spine, or TrPs in the sOMs upon palpation.

Course of measurement

This experimental study focused on determining the immediate effect of myofascial techniques on postural stability

in patients with reflex changes in the upper cervical region. During the initial clinical examination, postural stability was assessed using a stabilometric platform (Freestep Software; Sensor Medica, Rome, Italy). The platform's overall surface area measures 635 mm × 700 mm, while the active sensor surface spans 500 mm × 600 mm. Real-time sampling occurs at a frequency range of 300–350 Hz. Subsequently, the patients received MFR in the upper cervical region (Figure 1), as described in detail in the study by Ramezani et al. [25]. The total treatment duration was up to 10 min, and immediately after the treatment, all patients, as well as participants in the control group, underwent repeated posturographic examination. The examination was then repeated for all participants after an hour of rest.



Figure 1. Myofascial release technique in the upper cervical region

During the stabilometric assessment, body oscillations were evaluated by placing the subject on a force platform (Sensor Medica, Rome, Italy) in an upright position with arms at the sides, heels approximately 4 cm apart, and the forefoot rotated outward at an angle of 30°. This assessment was performed using the cervical test (Freestep software; Sensor Medica, Rome, Italy), which involves (1) standing with an upright head posture and eyes closed for 51.2 s, in accordance with the Romberg test, and (2) standing with head retroflexion and eyes closed for 51.2 s (Figure 2). The head retroflexion



Figure 2. Cervical test with upright head posture with eyes closed and with head retroflexion

posture is explained by several authors as causing deafferentation of proprioceptors in the cranial region, which is crucial for maintaining postural stability [28]. Posturographic recordings with the patient's head in retroflexion have been proposed as a reliable and sensitive test for diagnosing stability related to cervical dysfunction and as a diagnostic “fine-tuning” of stabilometry [29].

Measured parameters and data processing

To evaluate postural stability, we used parameters that characterise the movement of the Center of Pressure (CoP) in a standing position (Figure 3). The displacement of CoP was analysed by assessing (1) CoP trajectory length, (2) CoP ellipse area, (3) mean CoP velocity, (4) mean CoP displacement in the anteroposterior direction, (5) mean CoP displacement in the mediolateral direction, (6) standard deviation of CoP movement in the mediolateral direction, and (7) standard deviation of CoP movement in the anteroposterior direction.

The data collected underwent processing using Statistica 13.0 software (TIBCO Software Inc., CA, USA). Given the limited number of subjects in the experimental group, we employed the non-parametric Wilcoxon rank-sum test to evaluate the significance of differences in the measured values. Differences were deemed significant at $\alpha = 0.05$.

Results

The immediate effect of manual treatment on reflex changes in the sOMs on postural stability was demonstrated in almost all assessed stabilometric parameters (Table 1). Under the condition of standing with an upright head posture and closed eyes, there was a statistically significant rise in postural instability following the therapeutic intervention, with significance noted at $p < 0.05$. The length of the CoP trajectory was 450 ± 127 before MFR treatment (pre) and 607 ± 168 after MFR treatment (post) ($p = 0.02$). The area of the CoP ellipse was 95 ± 112 before treatment (pre) and 218 ± 158 after treatment (post) ($p = 0.03$). The mean speed of CoP movement was 10 ± 2 before treatment (pre) and 11 ± 4 after treatment (post) ($p = 0.03$). The mean value of CoP movement in the anteroposterior direction was -14 ± 10 before treatment (pre) and -26 ± 9 after treatment (post) ($p = 0.04$). The standard deviation of CoP movement in the anteroposterior direction was 1.8 ± 0.7 before treatment (pre) and 2.3 ± 1 after treatment (post) ($p = 0.01$). In the lateral direction, it was 1.8 ± 0.8 before treatment (pre) and 4.4 ± 2 after treatment (post) ($p = 0.04$). After one hour of rest (post1), the assessed stabilometric parameters did not significantly differ from the pre-values in the standing position with an upright head posture.

In the control group, significant changes were only observed in the length of the CoP trajectory in an upright stance between pre (511 ± 210) and post (602 ± 162) measurements ($p = 0.03$) (Table 2). In the head-tilted stance, significant changes were also observed in the length of the CoP trajectory after treatment, which persisted even after one hour of rest [pre (450 ± 190) vs post (474 ± 217)], $p = 0.02$; post1 = 486 ± 168 , $p = 0.04$). Significant changes were also noted in the mean velocity of CoP shortly after treatment [pre (10 ± 3) vs post (10 ± 2), $p = 0.02$].




Figure 3. Example of CoP Oscillations in the Cervical Test (Eyes Closed) before, immediately after, and one hour after therapy in a patient with TrPs in the sOMs

Table1 Comparison of stabilometric in the experimental group

Evaluated parameters	Pre (mean ± SD)	Post (mean ± SD)	Post1 (mean ± SD)	Pre vs. post		Pre vs. post1	
				Z	p	Z	p
Upright stance							
CoP trajectory length (mm)	450 ± 127	607 ± 168	494 ± 137	−2.1	0.02*	−1.1	0.15
CoP ellipse area (mm²)	95 ± 112	218 ± 158	108 ± 59	−1.9	0.03*	−0.8	0.21
Average CoP velocity (mm/s)	10 ± 2	11 ± 4	10 ± 2	−1.8	0.03*	−0.9	0.19
CoP X mean (mm)	2 ± 10	2 ± 8	2 ± 10	−0.2	0.42	−0.1	0.45
CoP Y mean (mm)	−14 ± 10	−26 ± 9	−21 ± 14	−1.7	0.04*	−1.5	0.07
CoP X SD (mm)	1.8 ± 0.7	2.3 ± 1	1.7 ± 0.7	−2.3	0.01*	−1.9	0.03*
CoP Y SD (mm)	1.8 ± 0.8	4.4 ± 2	3 ± 1.3	−1.7	0.04*	−1	
Head tilted stance							
CoP trajectory length (mm)	490 ± 142	518 ± 131	507 ± 164	−0.5	0.31	−1	0.17
CoP ellipse area (mm²)	103 ± 64	131 ± 111	95 ± 85	−0.3	0.39	−1.1	0.13
Average CoP velocity (mm/s)	10 ± 3	11 ± 3	10 ± 3	−0.1	0.44	−0.7	0.26
CoP X mean (mm)	1 ± 7	2 ± 6	3 ± 7	−0.3	0.36	−1.2	0.11
CoP Y mean (mm)	−18 ± 11	−17 ± 13	−17 ± 13	−0.3	0.39	−0.1	0.45
CoP X SD (mm)	1.7 ± 0.7	1.8 ± 0.7	1.6 ± 0.8	−0.1	0.44	−0.3	0.36
CoP Y SD (mm)	2.9 ± 1.3	3.4 ± 1.9	2.7 ± 1.3	−0.4	0.36	−1.1	0.14

Pre – before treatment, Post – after treatment, Post1 – one hour after treatment, CoP X mean – mean value of CoP movement in the anteroposterior direction, CoP Y mean – mean value of CoP movement in the mediolateral direction, CoP Y SD – standard deviation of CoP movement in the mediolateral direction, CoP X SD – standard deviation of CoP movement in the anteroposterior direction
* statistically significant value

Table 2 Comparison of stabilometric parameters in the control group

Eavaluated parameters	Pre (mean ± <i>SD</i>)	Post (mean ± <i>SD</i>)	Post1 (mean ± <i>SD</i>)	Pre vs. post		Pre vs. post1	
				<i>Z</i>	<i>p</i>	<i>Z</i>	<i>p</i>
Upright stance							
CoP trajectory length (mm)	511 ± 210	602 ± 162	551 ± 147	−1.9	0.03*	−1.2	0.12
CoP ellipse area (mm²)	83 ± 79	92 ± 61	86 ± 63	−0.1	0.49	−0.1	0.48
Average CoP velocity (mm/s)	10 ± 4	12 ± 3	11 ± 3	−1.4	0.08	−0.6	0.26
CoP X mean (mm)	1 ± 8	3 ± 6	6 ± 9	−0.4	0.36	−1.2	0.10
CoP Y mean (mm)	−20 ± 15	−22 ± 13	−20 ± 13	−0.4	0.34	−0.2	0.42
CoP X <i>SD</i> (mm)	1.9 ± 0.8	1.7 ± 0.9	1.9 ± 0.8	−0.7	0.24	−0.2	0.43
CoP Y <i>SD</i> (mm)	2 ± 1.3	2.8 ± 0.9	2 ± 1	−0.9	0.18	−0.3	0.37
Head tilted stance							
CoP trajectory length (mm)	450 ± 190	474 ± 217	486 ± 168	−2.1	0.02*	−1.8	0.04*
CoP ellipse area (mm²)	57 ± 40	45 ± 26	57 ± 38	−0.5	0.30	−1.6	0.05
Average CoP velocity (mm/s)	10 ± 3	10 ± 2	10 ± 3	−2.05	0.02*	−1.2	0.12
CoP X mean (mm)	3 ± 6	3 ± 5	4 ± 8	−0.7	0.23	−0.2	0.41
CoP Y mean (mm)	−18 ± 13	−20 ± 13	−21 ± 10	−0.6	0.26	−0.7	0.25
CoP X <i>SD</i> (mm)	1.8 ± 1	1.3 ± 0.5	1.5 ± 0.7	−0.5	0.29	−0.9	1.76
CoP Y <i>SD</i> (mm)	1.6 ± 0.7	1.6 ± 0.6	1.9 ± 0.8	−0.8	0.22	−1.3	0.10

Pre – before treatment, Post – after treatment, Post1 – one hour after treatment, CoP X mean – mean value of CoP movement in the anteroposterior direction, CoP Y mean – mean value of CoP movement in the mediolateral direction, CoP Y SD – standard deviation of CoP movement in the mediolateral direction, CoP X SD – standard deviation of CoP movement in the anteroposterior direction
* statistically significant value

Discussion

The objective of this study was to investigate whether the MFR technique would have an effect on stabilometric parameters in patients with TrPs in the sOMs. Surprisingly, the results indicated a worsening of certain stabilometric parameters immediately after the MFR treatment.

While many studies have focused on the effects of myofascial techniques and other manual therapy approaches on TrPs in the sOMs, they mostly describe subjective improvements such as pain relief or increased range of motion in the cervical joints [7, 18, 23]. However, the impact on postural stability has not been explored. It can be assumed that TrPs in the sOMs significantly influence postural stability because the upper part of the cervical spine plays a crucial role as a communication centre for mediated and coordinated reflexes that help maintain body position. Therefore, any intervention in these muscles can elicit an unexpected reaction from the postural system.

Although these findings contradict the results of several studies examining the impact of manual therapy on postural control, we did not find any studies investigating postural stability immediately after sOM therapy, nor any studies examining the effects of any manual technique on sOMs in relation to postural stability. A similar study by Fisher et al. [30] examined the immediate impact of cervical spine manipulation on postural stability, but their study failed to demonstrate any influence on CoP parameters. Additionally, Amato et al. [12] demonstrated improvements in some stabilometric indicators after self-treatment with the MFR technique in patients with migraines following 10 days of therapy. Similarly, Reid et al. [31] reported improved postural stability in a head-tilted standing position in patients with dizziness and neck pain after mobilisation techniques in the cervical area, but the improvements persisted after several weeks of observation.

Some authors have examined the effects of manipulative techniques on the cervical spine in asymptomatic individuals and found improved postural control after a single manipulation [14, 32]. As for the treatment specifically involving manual therapy in the cervical spine and its impact on postural stability, several explanations have been presented in the literature to account for the results of our study. However, all of these studies investigated MFR on postural stability after an extended period following the intervention (days to weeks).

The most significant factor associated with CoP deviations in patients with symptoms indicating cervical spine disorders is pain. Schabrun et al. [33] illustrated alterations in the somatosensory cortex during and post-pain, whereas modifications in the motor cortex occurred solely after pain cessation, exerting a more pronounced effect on CoP deviation. All patients in our study reported neck pain, which may have influenced the results. Moreover, patients with TrPs in the sOMs also reported head pain when placing their heads on a pillow, which can be explained by the effect of the weight of the occipital bone on TrPs in the suboccipital region [6].

Abnormal cortical processing of proprioceptive information during neck pain and its impact on postural stability has been described by several authors [34, 35] and supported by theories suggesting that muscle spindles increase their resting excitability by up to 200 % after muscle stretching compared to pre-stretch conditions [36]. Although the MFR technique does not directly stretch the muscles, it still leads to increased muscle spindle response and delayed muscle relaxation after muscle contraction [37]. Despite these theories, Peng et al. [3] point out in their narrative review on impaired proprioception in neck pain that high-quality research in this

area is still lacking. Passive head movements, which are part of manual therapy, as well as changes in position from lying to standing, may also impact increased CoP deviations in patients with long-standing cervical spine difficulties [30, 33]. Burstein et al. [7] also mention an important factor related to changes in CoP deviations, namely that muscle relaxation in the cervical spine area through MFR causes a sudden vasodilatory effect. Another theory worth mentioning is “sensory weighting”, which suggests that the postural and central nervous systems can reorganise the hierarchy of sensory information and prioritise other systems when the quality of sensory information is compromised, such as the vestibular system or proprioception from the lower limbs. [38, 39]. Hypothetically, such sensory weighting could have occurred in the patients in this study, and after intervention with TrPs in the sOM, it could have led to a new reorganisation of sensory information, potentially resulting in larger CoP deviations.

Limitations

The primary constraint in our study was the limited sample size, which was a result of excluding discrete subjects. Assessing stability parameters at earlier time intervals following MFR treatment would provide a more precise understanding of when stability adjustments occur and give more accurate insights into the timing of stability adjustments after MFR treatment.

Conclusions

In this study on the influence of the MFR technique on postural stability immediately after its application, we observed significant differences in posturographic parameters, indicating increased instability in patients compared to their pre-therapy state. The patients received myofascial treatment for TrPs in the sOMs. However, after an hour of rest, the patients' postural parameters were adjusted and normalised. The myofascial technique also had an impact on healthy individuals in the control group, particularly on the trajectory of the CoP. Overall, the influence of MFR on postural stability was much smaller in healthy individuals without TrPs in the suboccipital region. These results support the theory that therapeutic intervention in the upper cervical spine affects proprioception from that area and postural stability.

For safety reasons, we recommend that patients remain in a stable position that does not require significant postural stability after therapy. This is particularly important for patients scheduled to undergo additional kinesiotherapy after manual therapy, as it could affect safety and the quality of exercise. These findings could contribute to improving the safety and quality of rehabilitation treatment for patients.

Ethical approval

The research related to human use has complied with all the relevant national regulations and institutional policies, has followed the tenets of the Declaration of Helsinki, and has been approved by the Faculty of Health Care, University of Prešov (approval No.: EK-FZOPU 34/17.3.2023).

Informed consent

Informed consent has been obtained from all individuals included in this study.

Disclosure statement

No author has any financial interest or received any financial benefit from this research.

Conflict of interest

The authors state no conflict of interest.

Funding

This research received no external funding.

References

- [1] Mc Caskey MA, Schuster-Amft C, Wirth B, Suica Z, de Bruin ED. Effects of proprioceptive exercises on pain and function in chronic neck and low back pain rehabilitation: a systematic literature review. *BMC Musculoskelet Disord*. 2014;15:382; doi: 10.1186/1471-2474-15-382.
- [2] McPartland JM, Brodeur RR. Rectus capitis posterior minor: a small but important suboccipital muscle. *J Bodyw Mov Ther*. 1999;3(1):30–5; doi: 10.1016/S1360-8592(99)80040-0.
- [3] Peng B, Yang L, Li Y, Liu T. Cervical proprioception impairment in neck pain-pathophysiology, clinical evaluation, and management: a narrative review. *Pain Ther*. 2021;10(1):143–64; doi: 10.1007/s40122-020-00230-z.
- [4] Humphreys BK. Cervical outcome measures: testing for postural stability and balance. *J Manipulative Physiol Ther*. 2008;31(7):540–6; doi: 10.1016/j.jmpt.2008.08.007.
- [5] Mergner T, Nasios G, Maurer C, Becker W. Visual object localisation in space. Interaction of retinal, eye position, vestibular and neck proprioceptive information. *Exp Brain Res*. 2001;141(1):33–51; doi: 10.1007/s002210100826.
- [6] Simons DG. Understanding effective treatments of myofascial trigger points. *J Body Work Mov Ther*. 2002;6: 81–8; doi: 10.1054/jbmt.2002.0271.
- [7] Burstein R, Nosedá R, Borsook D. Migraine: multiple processes, complex pathophysiology. *J Neurosci*. 2015;35 (17):6619–29; doi: 10.1523/JNEUROSCI.0373-15.2015.
- [8] Fernández de las Peñas C, Alonso-Blanco C, Cuadrado ML, Gerrwin R, Pareja J. Myofascial trigger points in the suboccipital muscles and forward head posture in chronic tension type headache. *Headache*. 2006;46: 454–60; doi: 10.1111/j.1526-4610.2006.00288.x.
- [9] Ciubotariu A, Arendt-Nielsen L, Graven-Nielsen T. The influence of muscle pain and fatigue on the activity of synergistic muscles of the leg. *Eur J Appl Physiol*. 2004; 91(5–6):604; doi: 10.1007/s00421-003-1026-9.
- [10] Thunberg J, Ljubisavljevic M, Djupsjobacka M, Johansson H. Effects on the fusimotor-muscle spindle system induced by intramuscular injections of hypertonic saline. *Exp Brain Res*. 2002;142(3):319–26; doi: 10.1007/s00221-001-0941-4.
- [11] Yang L, Yang C, Pang X, Li D, Yang H, Zhang X, Yang Y, Peng B. Mechanoreceptors in diseased cervical intervertebral disc and vertigo. *Spine*. 2017;42(8):540–6; doi: 10.1097/BRS.0000000000001801.
- [12] Amato A, Messina G, Giustino V, Brusa J, Brighina F, Proia P. A pilot study on non-invasive treatment of migraine: The self-myofascial release. *Eur J Transl Myol*. 2021;31(1):9646; doi: 10.4081/ejtm.2021.9646.
- [13] Jull G, Amiri M, Bullock-Saxton J, Darnell R, Lander C. Cervical musculoskeletal impairment in frequent intermittent headache. Part 1: subjects with single headaches. *Cephalalgia*. 2007;27(7):793–802; doi: 10.1111/j.1468-2982.2007.01345.x.
- [14] Smith L, Mehta M. The effects of upper cervical complex high velocity low amplitude thrust technique and sub-occipital muscle group inhibition techniques on standing balance. *J Osteop Med*. 2008;11:149–68; doi: 10.1016/j.ijosm.2008.08.020.
- [15] Fernández de las Peñas C, Cuadrado ML. Therapeutic options for cervicogenic headache. *Expert Rev Neurother*. 2014;14(1):39–49; doi: 10.1586/14737175.2014.863710.
- [16] Karegeannes M, Fernández De las Peñas C, Finnegan M. Suboccipital muscles. In: Travell JG, Simons DG, Simons LS (eds). *Myofascial Pain and Dysfunction*. 3rd ed. Philadelphia: Wolters Kluwer; 2019, pp. 178–84.
- [17] Treleaven J. Sensorimotor disturbances in neck disorders affecting postural stability, head and eyemovement control. *Man Ther*. 2008;13(1):2–11; doi: 10.1016/j.math.2007.06.003.
- [18] Moraska A, Chandler C. Changes in clinical parameters in patients with tension-type headache following massage therapy: a pilot study. *J Man Manip Ther*. 2008; 16(2):106–12; doi: 10.1179/106698108790818468.
- [19] Giacomini P, Di Maure R, Topaxio D, Girolamo SD. Head tilt posturography: clinical value in peripheral labyrinthine disorders. *Ann Otol Rhinol*. 2015;2(12):1073.
- [20] Sprenger A, Wojak JF, Jandl NM, Helmchen CH. Postural control in bilateral vestibular failure: Its relation to visual, proprioceptive, vestibular, and cognitive input. *Front Neurol*. 2017;8:444; doi: 10.3389/fneur.2017.00444.
- [21] Jull G, Stanton W. Predictors of responsiveness to physiotherapy management of cervicogenic headache. *Cephalalgia*. 2005;25(2):101–8; doi: 10.1111/j.1468-2982.2004.00811.x.
- [22] Dommerholt J, Bron C, Franssen J. Myofascial trigger points: an evidence-informed review. *J Man Manipul Ther*. 2006;14(4):203–21; doi: 10.1179/106698106790819991.
- [23] Chaitow L, Delany J. Neuromuscular techniques in orthopedics. *Tech Orthop*. 2003;18(1):74–86; doi: 10.1097/00013611-200303000-00013.
- [24] Kamonseki DH, Lopes EP, Meer HA, Calixtre LB. Effectiveness of manual therapy in patients with tension-type headache. *Asystematic review and meta-analysis*. *Disabil Rehabil*. 2020;12:1–10; doi: 10.1080/09638288.2020.1813817.
- [25] Ramezani E, Arab AM, Nourbakhsh MR. Sub-occipital myofascial release technique in subjects with cervicogenic headache. *Pharmacophore*. 2017;8(6S):e-1173299; doi: 10.1016/j.jbmt.2018.09.041.
- [26] Schleip R. Fascial plasticity – a new neurobiological explanation Part 2. *J Bodyw Mov Ther*. 2003;7(2):104–16; doi: 10.1016/S1360-8592(02)00076-1.
- [27] Beinert K, Taube W. The effect of balance training on cervical sensorimotor function and neck pain. *J Mot Behav*. 2013;45(3):271–8; doi: 10.1080/00222895.2013.785928.
- [28] Norré ME. Posturography: head stabilisation compared with platform recording. Application in vestibular disorders. *Acta Otolaryngol Suppl*. 1995;520(2):434–6; doi: 10.3109/00016489509125291.
- [29] Nacci A, Ferrazzi M, Berrettini S, Panicucci E, Matteucci J, Bruschini L, Ursino F, Fattori B. Vestibular and stabilometric findings in whiplash injury and minor head trauma. *Acta Otorhinolaryngol Ital*. 2011;31:378–89.
- [30] Fisher AR, Bacon CJ, Mannion JVH. The effect of cervical spine manipulation on the postural sway in patients with non-specific neck pain. *J Manipul Physiol Ther*. 2015;38(1):65–73; doi: 10.1016/j.jmpt.2014.10.014.
- [31] Reid SA, Rivett DA, Katekar MG, Callister R. Sustained natural apophyseal glides (SNAGs) are an effective treatment for cervicogenic dizziness. *Man Ther*. 2008;13(4): 357–66; doi: 10.1016/j.math.2007.03.006.
- [32] Nolan JH. The Effect of cervical spine chiropractic manipulation on balance. Johannesburg: Department of Chiropractic, University of Johannesburg; 2009.

- [33] Schabrun SM, Jones E, Kloster J, Hodges PW. Temporal association between changes in primary sensory cortex and corticomotor output during muscle pain. *Neurosci.* 2013;235(3):159–64; doi: 10.1016/j.neuroscience.2012.12.072.
- [34] Brumagne S, Janssens L, Janssens E, Goddyn L. Altered postural control in anticipation of postural instability in persons with recurrent low back pain. *Gait Posture.* 2008; 28(4):657–62; doi: 10.1016/j.gaitpost.2008.04.015.
- [35] Paulus I, Brumagne S. Altered interpretation of neck proprioceptive signals in persons with subclinical recurrent neck pain. *J Rehabil Med.* 2008;40(6):426–32; doi: 10.2340/16501977-0189.
- [36] Pickar JG, Bolton PS. Spinal manipulative therapy and somatosensory activation. *J Electromyogr Kinesiol.* 2012; 22(5):785–94; doi: 10.1016/j.jelekin.2012.01.015.
- [37] Schomacher J, Farina D, Lindstroem R, Falla D. Chronic trauma-induced neck pain impairs the neural control of the deep semispinalis cervicis muscle. *Clin Neurophysiol.* 2012;123(7):1403–8; doi: 10.1016/j.clinph.2011.11.033.
- [38] Gajdoš M, Nechvátal P, Kendrová L, Čuj J, Kozel M. The effect of myofascial therapy of the suboccipital muscles on postural stability – a pilot study. *Rehabil Fyz Lek.* 2022; 29(4):208–13; doi: 10.48095/ccrhfl2022208.
- [39] Vuillerme N, Pinsault N, Vaillant J. Postural control during quiet standing following cervical muscular fatigue: effects of changes in sensory inputs. *Neurosci Lett.* 2005; 378(3):135–9; doi: 10.1016/j.neulet.2004.12.024.