

Exploring the impact of physical, cardiorespiratory, and functional indicators on health-related quality of life in active and healthy older adults: a cross-sectional study

DOI: <https://doi.org/10.5114/pq/207970>

Anna Arnal-Gómez¹ , Elena Muñoz-Gómez² , Pedro Lizondo-Osset² , Pablo Sampietro-Picó² ,
Maryana Babetska² , Gemma V. Espí-López³ 

¹ Physiotherapy in Motion, Multi-speciality Research Group (PTinMOTION), University of Valencia, Valencia, Spain

² Department of Physiotherapy, Faculty of Physiotherapy, University of Valencia, Valencia, Spain

³ Exercise Intervention for Health (EXINH), University of Valencia, Valencia, Spain

Abstract

Introduction. Population ageing is a growing social phenomenon with significant health implications. This study aimed to analyse health-related quality of life (QoL) and its relation to physical activity, musculoskeletal pain, balance, cardiorespiratory, and functional characteristics in community-dwelling older adults (> 60 years) who meet the criteria for successful ageing.

Methods. A cross-sectional study was conducted using a non-probabilistic sample of physically active older adults. Primary outcomes included physical and mental QoL (SF-12). Secondary outcomes were physical activity (IPAQ and Exercise Benefits/Barriers Scale), musculoskeletal pain (VAS), balance (Unipedal Stance Test, Y-Balance Test), cardiorespiratory variables (heart rate, oxygen saturation, and dynamic lung volumes via spirometry), and functional capacity (Six-Minute Walk Test and Timed Up and Go test). Shapiro–Wilk tested normality, and Levene’s test assessed homoscedasticity. Pearson correlations explored associations between QoL and secondary outcomes (i.e., physical activity, musculoskeletal pain, dynamic lung volumes by forced spirometry, and functional capacity). Stepwise linear regression was used to identify predictors of QoL (i.e., respiratory and functional factors).

Results. 91 participants (mean age = 69.20 ± 5.19 years; 70.33% female) were assessed. Moderate physical activity correlated with both physical ($r = 0.25, p < 0.05$) and mental ($r = 0.27, p < 0.05$) QoL. Physical QoL was associated with pain and forced vital capacity ($R^2 = 0.26; F(2,84) = 14.68, p < 0.01$), while mental QoL was associated with physical QoL and Timed Up and Go performance ($R^2 = 0.19, F(2,84) = 9.89, p < 0.01$).

Conclusions. In active older adults, physical QoL was associated with pain and lung function, while mental QoL was linked to physical QoL and functional mobility. These factors should be prioritised in health strategies aimed at promoting successful ageing.

Key words: dry needling, exercise, functional status, range of motion, pain

Introduction

Currently, the population is ageing significantly, with the fastest increase projected for the ‘oldest’ old, and this has become a social phenomenon that will lead to changes both in older adults’ health and in society as a whole [1]. Considering that ageing has been related to physical, cardiorespiratory, and functional decline, and that there is a growing number of older adults reaching their 90s and 100s [2], the challenge is how to expand both the healthy and functional years in the life span.

Recently, the concept of successful ageing, based on a low probability of suffering illness or disability, great physical, functional and mental capacity, and an active commitment to life, has been studied [3]. Older adults who age successfully can maintain their functionality for longer and have lower morbidity and disability [4]. Therefore, it has become of paramount importance to understand and encourage successful ageing in older adults.

Moreover, it is crucial to focus on quality of life (QoL) in older adult groups, since it can help identify issues that are frequently linked to dysfunction, whether in the physical or mental domain [5]. For older adults, the concept of health-related QoL encompasses various dimensions, including physi-

cal health, mental state, levels of functionality, social interaction and context [6].

Physical activity can be a factor that significantly differentiates the average level of QoL for both middle-aged and older adults [7]. The level of physical activity also seems to play an important role, since a high practice of physical activity has shown to favour a functional capacity that can, in turn, guarantee the autonomy and independence in activities of daily life [8]. Moreover, physical activity can help older adults preserve cognitive function, delay dependency, and reverse frailty [9, 10]. Recently, a review has highlighted that walking can lead to improved lung function in healthy older adults [11]. In fact, inactive behaviour and sedentary lifestyles increase with age and are major risk factors for chronic health conditions [12].

In spite of this evidence, previous reports have shown that only 25% of individuals over 65 meet the recommended minimum activity levels for maintaining health [13]. This might be attributed to the impact of the ageing process on the health parameters of older adults. Firstly, it is well-documented that the decline in the musculoskeletal system’s capacities can result in weakness, frailty, reduced balance, diminished strength, and impaired functionality, among other issues [14]. Moreover, older adults have an increased inci-

Correspondence address: Elena Muñoz-Gómez, Department of Physical Therapy, University of Valencia, Gascó Oliag Street nº 5, 46010 Valencia, Spain, e-mail: elena.munoz-gomez@uv.es; <https://orcid.org/0000-0001-9137-1656>

Received: 18.11.2024

Accepted: 07.07.2025

Citation: Arnal-Gómez A, Muñoz-Gómez E, Lizondo-Osset P, Sampietro-Picó P, Babetska M, Espí-López GV. Exploring the impact of physical, cardiorespiratory, and functional indicators on health-related quality of life in active and healthy older adults: a cross-sectional study. *Physiother Quart.* 2026;34(1):83–91; doi: <https://doi.org/10.5114/pq/207970>.

dence of painful conditions related to musculoskeletal, nervous system, or metabolic changes, with studies reporting between 25–76% of community-dwelling older adults suffering some type of pain [15]. Common musculoskeletal conditions in older adults include neck pain, primary headaches, and shoulder discomfort [16]. Previous literature has stated that older adults might have different views and understandings of pain compared to younger adults, but the ideal source of pain history is the patient (self-report) [16].

These changes not only affect skeletal muscles but also the cardiorespiratory system undergoes a series of slow and progressive morphological and functional modifications with the ageing process [17]. All these changes and conditions have a strong impact on older people's health-related QoL, which has been shown to decline with advancing age and to be reduced among inactive older adults with diminished physical performance [18]. However, no previous study has focused on how being physically active and ageing parameters such as pain, balance, cardiorespiratory or functional ones can influence and explain how to have a successful QoL. Therefore, less focus has been given to comprehensively understanding factors related to QoL in successful and active older adults. All considered, physical, cardiorespiratory, and functional abilities, parameters that are all so intricately affected with ageing, could be enhanced and have, in turn, a strong influence on achieving a successful and active ageing, and could make a profound effect on older adults' health and QoL. Thus, to explore complex relationships and make forecasts by examining how these multiple factors influence QoL, a multiple regression analysis should be applied. The hypothesis of this study was that successful older adults who are physically active will have positive health parameters, and these will influence their QoL.

Therefore, the aim of this study was to describe and analyse QoL, physical activity, musculoskeletal pain, balance, cardiorespiratory and functional characteristics of older adults over 60 years of age, who comply with the concept of successful ageing. It also aimed to analyse the possible associations between physical and mental health-related QoL and physical, cardiorespiratory and functional health indicators.

Subjects and methods

Design

A cross-sectional study was performed at the Laboratories of the University of Valencia (Spain) from February to April 2021. Participants were assessed once by a physiotherapist. The manuscript was written in accordance with the Strengthening the Reporting of Observational studies in Epidemiology (STROBE) protocol [20].

Participants

Participants were recruited from older adults' associations of Valencia (Spain). A non-probability sampling approach was used to select the study participants. Inclusion criteria were: community-dwelling older adults (considered as those of over 60 years of age [21]); with successful ageing, defined as those who had not suffered stroke, coronary artery disease, angina, acute myocardial infarction, cancer, or diabetes; plus they had optimal cognitive, physical, mental, respiratory and vascular function and a lack of disability while being functionally independent [22]; and considered as active by the International Physical Activity Questionnaire (IPAQ) short version [23]. Exclusion criteria were: Mini-Mental State Exami-

nation < 25 points (administered by a trained interviewer) and suffering any acute injury (acute musculoskeletal or osteoarticular pain).

Outcomes

All participants undertook a face-to-face assessment performed by a physiotherapist with more than 10 years of experience that included:

Anthropometric variables: age, sex, body weight (kg) measured by a Tanita BC 601 weighing device (TANITA Ltd., Tokyo, Japan), and height (cm) measured with a stadiometer SECA 213 (Seca Ltd., Hamburg, Germany), with both instruments having been shown to be reliable and valid for use in community assessments [24]. Also, body mass index (BMI) was calculated (kg/m^2).

Primary outcome

Quality of life: The Short Form 12 (SF-12) Health Questionnaire v2 was used, which measures the impact of health on everyday life. It consists of 12 items that include areas related to physical and mental aspects of health, thus determining a physical subscale and a mental subscale using data from the Medical Expenditure Panel Survey longitudinal panel (2014–2015). The maximum score that can be obtained is 100 points, which indicates the highest level of health. It is a reliable and valid instrument with high internal consistency for both physical and mental components (Cronbach's alpha: 0.87 and 0.86, respectively) and good and moderate test-retest validity (intraclass correlation coefficient (ICC): 0.79 and 0.59, respectively) [25].

Secondary outcomes

Physical activity variables: (1) Level of physical activity, calculated by the number of METs/week, and then classified accordingly (moderate-intensity, level 2, and high-intensity, level 3) by the International Physical Activity Questionnaire (IPAQ) short version [23]; (2) Number of days per week participants exercised; (3) Type of exercise; (4) Perception of exercise, benefits and barriers, using the Exercise Benefits/Barriers Scale (EBBS), composed of 43 items evaluated using a 4-point Likert scale ('completely agree' to 'completely disagree'); the higher the score, the more positive the perception of exercise. The reliability coefficient (Cronbach's alpha) is 0.95 for the total scale [26].

Musculoskeletal pain: patients were asked if they felt musculoskeletal pain and to rate the intensity of this musculoskeletal pain intensity with the Visual Analogue Scale (VAS). The VAS is a unidimensional measure of non-specific perceived pain intensity that is appropriate for use with older adults [27]. It is a continuous scale composed of a horizontal line typically 10 cm in length, on which the participant is asked to indicate pain intensity with a score of 0 being 'no pain' and 10 being 'the worst pain imaginable', a higher score indicates greater pain intensity. Participants were asked to indicate the body area that was most painful and to rate the intensity of the 'current' pain or pain intensity 'in the last 24 hours' for that area, the VAS has a high internal consistency (0.97) [28]. Also, an ad hoc question was asked with a yes/no answer indicating the need of medication and/or visiting a physiotherapist for this pain.

Balance: (1) Static balance, measured by two variables: Static balance, with the Unipedal Stance Test (UPST) which consists of assessing the time the participants can maintain

eyes closed monopodal support while their arms are crossed at chest level and one leg is raised [29]. Three repetitions were done, and the best was selected for analysis. This test has shown an ICC of 0.998 for eyes closed [29]. (2) Dynamic balance was assessed with a variant of the Star Excursion Balance Test (SEBT), the Lower Quarter Y-Balance Test (YBT-LQ) which is a reliable and valid measure (ICCs for intratester reliability 0.78–0.96 and ICCs for intertester reliability were 0.35–0.84) [30]. Participants have to maintain stability while moving one lower limb in three directions (anterior, posterolateral and posteromedial) and keeping the other limb stable. For the analysis, the mean of three attempts in each direction was recorded [31]. A principal component analysis was performed to summarise the three directions of the right leg in one variable and the three of the left leg in another variable.

Cardiorespiratory variables: (1) Pulse oximetry: Oxygen saturation (SpO_2) and heart rate were assessed with a pulse oximeter, SmartOx (WEINMANN, Medical Technology, Hamburg, Germany). Oxygen saturation has shown to be a reliable method for detecting hypoxaemia and has been reported accurate in reflecting one-point measurements of SpO_2 . (2) Forced spirometry using a spirometer Spirovit SP-10 (Contec, Barcelona) was performed according to American Thoracic Society standards. Dynamic lung volumes such as forced vital capacity (FVC) and the forced expired volume at 1 second (FEV1) were obtained by the forced expiratory flow manoeuvre, and the FEV1/FVC ratio was calculated. For FVC and FEV1, an 80% or over is an optimal value [32], and for the FEV₁/FVC ratio, a decreased percentage defines the concept of 'obstruction'. Spirometry has shown an ICC of 0.89 in COPD patients [33].

Functional variables: (1) The Six-Minute Walk Test (6MWT), which aims to walk for 6 minutes around the 30 m distance between cones. Pulse oximetry, final heart rate, SpO_2 , and dyspnoea were recorded using the Borg scale. The 6MWT is a reliable measure (ICC = 0.82–0.99) [34]; (2) Timed Up and Go test (TUG), where participants had to stand up from a chair, walk a 3 m distance, turn around and walk back, and sit back on the chair. The test is repeated 3 times and then the mean is calculated (i.e. < 10 Independent Mobility, 10–20 Mostly Independent, 20–29 variable mobility, > 30 reduced mobility [35]). It has shown excellent interrater reliability and varied intrarater reliability [36].

Sample size calculation

G*Power version 3.1.9.7 was used to calculate the sample size for the linear multiple regression analysis. Five predictor variables were established, along with a medium effect size ($f^2 = 0.15$), a type I error (α) of 5%, and a type II error (β) of 20%, according to a previous study conducted by Wanderley et al. [37]. The findings suggested that 92 people would be needed for the sample.

Statistical analysis

Statistical analyses were carried out through the SPSS program (v. 25, SPSS Inc., Chicago, IL, USA). Normality of variables was assessed using the Shapiro–Wilk test. Levene's test assessed homoscedasticity. Data are shown as mean and standard deviation (SD) for continuous variables, and frequency and percentage for categorical ones. As the assumption of normality was fulfilled in the variables analysed, correlations between dependent and independent variables were analysed using the Pearson coefficient (0.10–0.29 values indicated a small effect, 0.30–0.49 moderate effect, and ≥ 0.50 large effect). The all-bivariate correlation was displayed

using a Pearson correlation heatmap in the Matlab software (Matlab 2023, Mathworks, Natick, MA, USA). The correlation heatmap is a graphical representation in which each variable is depicted by a row and a column, and the cells within the matrix display the correlations between them. The colour intensity in each cell is a key indicator of the magnitude and direction of the correlation. Darker hues indicate stronger correlations, while lighter colours indicate weaker correlations. Positive correlations (when one variable increases, the other variable tends to increase) are represented by a warm colour (red). Negative correlations (when one variable increases, the other variable tends to decrease) are represented by a cool colour (blue). The potential of respiratory and functional factors to predict the variability of physical and mental health was investigated using a multiple stepwise linear regression model. Coefficient (B), standardised beta coefficients (β), R^2 , adjusted R^2 , and p -values were used to assess the association's strength. Only the five independent variables with the highest Pearson correlation coefficient values were included in a prediction model, based on the results of the correlation test. The significance level was chosen at $p < 0.05$.

Results

Of the 121 initial participants, in the end, a total of 91 older adults met the inclusion criteria and were analysed. Thirty participants were excluded for not being willing to participate, not meeting the inclusion criteria or being injured prior to assessment. No adverse events were reported (Figure 1).

Participants had a mean (SD) age of 69.18 (5.08) years, with 70.33% being women. All participants exercised regularly, and nearly 80% performed this exercise with moderate intensity. Anthropometric and physical activity characteristics are shown in Table 1.

QoL, musculoskeletal pain, balance outcomes (i.e. UPST, YBT-LQ), cardiorespiratory outcomes (i.e. heart rate, SpO_2 , FVC, FEV₁, and FEV₁/FVC), and functional outcomes (i.e. 6MWT and TUG) are shown in Table 2. The 6MWT, heart rate, SpO_2 and dyspnoea parameters before the test were adequate and those registered after the test are shown. Regarding pain, 93.41% of the sample had musculoskeletal pain, and the pain intensity was 4.31 (SD = 2.36). Pain was distributed in different areas with the lumbar spine being the area where more participants had pain (8.13%), followed by head and/or neck pain (60.44%), knee pain (56.04%), dorsal spine (30.77%), shoulder pain (30.77%), hip and pelvic pain (29.67%) and ankle and/or foot pain (28.57%) (Supplementary material 1). A total of 32.25% took medication for the pain, and 77.50% regularly visited a physiotherapist with the aim of relieving pain.

Correlations and regression models

The Pearson Correlation Coefficient showed a statistically positive correlation between both physical and mental health-related QoL and higher levels of physical activity ($r = 0.25$, $p < 0.05$; and $r = 0.27$, $p < 0.05$, respectively) (Figure 2, and Supplementary material 2). Moreover, both physical and mental health-related QoL showed a statistically negative correlation with pain; so, the higher the pain, the lower the QoL. Physical health-related QoL showed a positive association with cardiorespiratory variables (SpO_2 , FVC, and FEV₁) static balance (UPST), functional parameters (6MWT distance and SpO_2 at the end of the test, and the TUG), and the mental health-related QoL. As for mental health-related QoL, it showed positive correlations with respiratory variables (FVC, and FEV₁), functional parameters (6MWT distance and the TUG), and physical health-related QoL.




Figure 1. Flow diagram of study participation

Table 1. Anthropometric and physical activity variables of the sample.

Variable	Total (n = 91)
Age [years, mean (SD)]	69.18 (5.09)
Sex [n (%)]	
women	64 (70.30%)
men	27 (29.70%)
BMI (kg/m ² , mean (SD))	25.69 (3.84)
IPAQ	
physical activity (METs/week, mean (SD))	2108.52 (1290.81)
moderate intensity [n (%)]	74 (79.12%)
high intensity [n (%)]	17 (18.68%)
Exercise frequency [n (%)]	
once a week	4 (4.40%)
2–3 times/week	38 (41.76%)
4–5 times/week	19 (20.88%)
every day	30 (33.97%)
Type of exercise [n (%)]	
walking	48 (52.75%)
hiking	12 (13.19%)
swimming	8 (8.79%)
therapeutic gymnastics	7 (7.69%)
yoga	2 (2.20%)
others	14 (15.38%)
EBBS (points) ^a	72.84 (21.49)

Descriptive analyses were performed.
 IPAQ – International Physical Activity Questionnaire,
 EBBS – Exercise Benefits/Barriers Scale

Table 2. Pain, balance, respiratory, and functional outcomes, and quality of life of the sample

Outcome	Total (n = 91) mean ± SD
Quality of life	
SF-12 physical subscale	73.07 (9.34)
SF-12 mental subscale	83.18 (11.65)
Musculoskeletal pain (VAS)	4.31 (2.36)
BALANCE	
UPST (s)	
right leg	3.39 (4.73)
left leg	2.87 (2.67)
YBT-LQ (cm)	
right leg	63.54 (12.00)
left leg	64.36 (13.32)
Cardio-respiratory variables	
HR (bpm)	80.57 (13.63)
SpO ₂	97.59 (1.37)
FVC (l)	2.75 (0.75)
FVC (%) predicted	82.18 (12.55)
FEV ₁ (l/min)	2.16 (0.69)
FEV ₁ (%) predicted	79.85 (18.12)
FEV ₁ /FVC (%)	78.18 (12.55)
Functional variables	
6MWT	
distance (m)	498.42 (62.33)
SpO ₂	96.81 (3.10)
HR (bpm)	92.23 (19.02)
dyspnoea	0.22 (0.56)
TUG (s)	6.72 (1.51)

Descriptive analyses were performed.
 VAS – Visual Analogue Scale, HR – heart rate, SpO₂ – saturation of peripheral oxygen, FVC (l) – forced vital capacity, FEV₁ – forced expiratory volume in one second, UPST – unipedal stance test, YBT-LQ – Lower Quarter Y-Balance Test, 6MWT – Six-Minute Walk Test, TUG – Timed Up and Go, SF-12 – Short Form 12 Health Survey




Figure 2. Correlation analyses were performed. Correlation heatmap between physical and mental health-related quality of life and the secondary outcomes (i.e., physical activity, musculoskeletal pain, balance, cardio-respiratory variables and functional variables). Darker colours indicate stronger correlations, while lighter colours indicate weaker correlations. Positive correlations (when one variable increases, the other variable tends to increase) are represented by a warm colour (red). Negative correlations (when one variable increases, the other variable tends to decrease) are represented by a cool colour (blue)

Table 3. Multiple stepwise linear regression analysis for quality of life in relation to mental and physical health

Dependent variable	Model	B	β	p -value	VIF	Overall model		
						R^2	F	SEE
Physical health (SF-12)	Predictor variables					0.26	14.68	8.12
	VAS	-1.50	-0.38	< 0.01*	1.04			
	FVC (l)	3.47	0.28	< 0.05*	1.04			
	excluded variables							
	FEV ₁ (l/min)		0.01	> 0.05	3.96			
Mental health (SF-12)	Predictor variables					0.19	9.89	10.66
	SF-12 physical subscale	0.38	0.30	< 0.01*	0.95			
	TUG	-1.99	-0.25	< 0.05*	0.95			
	excluded variables							
	FVC (l)		0.21	> 0.05	1.21			
FEV ₁ (l/min)		0.13	> 0.05	1.27				
physical activity (METs/week)		0.19	> 0.05	1.07				

Multiple stepwise linear regression analyses were performed.

B – regression coefficient, β – standardised coefficient, VIF – variance inflation factor, SEE – standard error of estimate, SF-12 – Short Form 12 Health Survey, VAS – Visual Analogue Scale, FVC – forced vital capacity, FEV₁ – forced expiratory volume in one second, TUG – Timed Up and Go * $p < 0.05$

For the regression analyses, only variables with significant correlations concerning physical health-related QoL and mental health-related QoL were considered (Table 3). For physical health-related QoL, the model showed an R^2 of 0.26, meaning that 25.90% of physical health variability was related to musculoskeletal pain and FVC ($F(2,84) = 14.68, p < 0.01$). For mental health, the model shows an R^2 of 0.19, indicating that 19.10% of mental health variability was related to physical health and TUG ($F(2,84) = 9.89, p < 0.01$).

Discussion

Our results showed that successful and active older adults tend to exercise with moderate physical activity, with walking 2–3 times a week being the most common physical activity. Globally, their mental health-related QoL was higher

than the physical, and 93.41% of the sample had musculoskeletal pain. Their cardiorespiratory parameters are adequate, and they have optimal functional independence. Their physical health-related QoL is significantly associated with musculoskeletal pain and cardiorespiratory parameters such as FVC. Mental health-related QoL is significantly related and can be predicted by adequate physical health-related QoL and good performance in the TUG functional parameter. Therefore, the overall QoL of successful and active older adults can be influenced by their respiratory conditions and their functional capacity; thus, these are areas healthcare professionals should focus on to improve QoL in the older population.

New disease patterns of older people have risen due to sedentary lifestyles, a lack of physical activity, stress, poor diets and consumption of toxic substances [38]. Compre-

hensively investigating, like in this study, which parameters favour a good QoL in people with successful ageing, allows clinicians to take them into account in practice, which could reduce the costs caused by ageing in primary care consultations as well as in the health system in general, both in terms of physical and psychological repercussions.

Physical health-related quality of life

Our results showed that physical health-related QoL is significantly associated with optimal levels of FVC. This may be explained because ageing is characterised by a reduction in muscle strength, joint mobility and cardiorespiratory capacity, and this reduction in physical capacities translates into greater breathing effort and a decrease in respiratory values. Moreover, there is a decline in lung function FVC, FEV₁, and peak expiratory flow rate as age increases [39]. Our participants, by having a successful ageing process and an active life, showed optimal spirometric results [40], with mean FVC and FEV₁ very close to 80% (82.18 and 79.85%, respectively) and a FEV₁/FVC of 78.18%. Our results highlight the importance of monitoring these variables in older adults, since they have shown a relationship with physical health-related QoL, which is ultimately one of the main objectives of geriatric healthcare. Furthermore, respiratory symptoms, frailty, and mortality are strongly associated with respiratory impairment [41]. Since respiratory parameters are feasible, simple and cost-effective to monitor through spirometry [32], it is important for health professionals to make use of these tools, so that preventive and treatment measures can be taken to achieve or maintain optimal levels. It should be noted that physical activity is a viable strategy for improving cardiorespiratory function [42], which also provides benefits in different physical, social and psychological dimensions [43]. Thus, promoting physical activity will improve their cardiorespiratory function, which in turn will enhance their physical health-related QoL.

In addition, our results showed that pain is another predictor of physical health; thus, higher levels of pain intensity tend to be associated with lower QoL. Chronic pain can have a significant impact on physical function, emotional well-being, and overall enjoyment of life, particularly in older adults, who may have other age-related health problems [44]. Therefore, it is important to manage pain effectively through individually tailored therapeutic approaches such as physiotherapy.

Mental health-related quality of life

Regarding mental health, our study has corroborated the relationship between physical and mental health-related QoL. According to the World Health Organization, more than 20% of people over 60 years old suffer from some type of mental disorder, with depression and dementia being the most prevalent. 3.8% of the older population is affected by anxiety disorders [45], commonly treated with antidepressants, but these may produce cardiovascular problems, epilepsy and an increased risk of mortality [46]. This makes it necessary to look for other alternatives, such as being active and practicing regular physical activity. Our results showed that functional capacity in terms of the TUG is associated with mental health. This is consistent with previous studies [47], which found that older adults with poorer mental health presented poorer performance on the TUG. In clinical practice, it is very interesting to be able to monitor functional capacity using the TUG, as it is a simple tool to use and offers vast information [48]. This in turn will make it possible to develop individualised prevention and treatment strategies for older adults.

Moreover, poor endurance and energy have an effect of 13.4% on the mental-health QoL [49]. In contrast, older persons who engage in more physical activity and have greater functional ability may be more resilient to anxiety and depression [50], and those with better functional capacity usually present higher vitality [37]. Interestingly, people with poor dynamic balance, functional mobility and gait as well as depressive symptoms often have a fear of falling [51], which may lead to a sedentary lifestyle. Our results also showed some correlations between balance variables and physical-health QoL; however, they were not significant in the predictive model, probably due to the need for a larger sample.

Future research directions

Our results, together with previous studies that highlight that exercise programs for older adults can reduce healthcare costs and prevent falls, strokes, heart attacks, and chronic diseases [10], enhance the idea of promoting research related to physical activity in older adults. Moreover, ageing is a complex process that must be conceived as a whole, which is in line with showing a relation between physical and mental health, so a multicomponent approach should be taken into consideration in future research.

Applications of the findings

Overall, improving respiratory, physical and functional variables may favour an active and successful lifestyle in older adults and enhance mental and physical health-related QoL. These findings are of interest for identifying risks and protective factors that favour successful ageing and developing strategies not only for prevention and treatment but also for promoting successful and active ageing in society.

Limitations

The sample consisted mainly of women, either because this is a situation that is a common characteristic to older adults of the population in Spain, or because women were more willing to collaborate. Thus, greater equality regarding sex would be important in future research. Additionally, although the sample size calculation indicated the need for 92 participants, only 91 individuals were ultimately included, which may slightly reduce the statistical power of the analyses and should be considered when interpreting the results. Future research would also benefit from a bigger sample size or even from analysing gender differences.

As strengths, this study includes a set of variables in active older adults with successful ageing that have not been previously studied or contrasted in relation to their impact on QoL. Therefore, this study confirms the common idea that physical activity is related to health parameters relevant to an optimal QoL.

Conclusions

Older adults with successful ageing who perform physical activity have optimal health parameters, and these parameters show relationships between them. Respiratory conditions and functional capacity have shown to influence the physical and mental health-related QoL of successful active older adults. Thus, these are areas healthcare professionals should focus on to improve QoL in the older population.

Database availability

The datasets generated and/or analysed during the current study are available from the corresponding author on reasonable request.

Acknowledgements

We gratefully acknowledge the participation of those who volunteered from the 'Amics de la Nau Gran', 'Hogares compartidos' and 'Centro de jubilados Salvador Allende' associations.

Ethical approval

The research related to human use complied with all the relevant national regulations and institutional policies, followed the tenets of the Declaration of Helsinki, and was approved by the Ethics Committee for Human Research of the University of Valencia, Spain (approval No.: H1539944398481), with ClinicalTrials.gov registration ID: NCT04732078, registered the 29th of January 2021.

The manuscript was written in accordance with the STROBE protocol.

Informed consent

Informed consent was obtained from all individuals included in this study.

Disclosure statement

No author has any financial interest or received any financial benefit from this research.

Conflict of interest

The authors state no conflict of interest.

Funding

This research received no external funding.

References

- [1] Alvarado-García A, Salazar-Maya A. Aging concept analysis. *Gerokomos*. 2014;25(2):57–62; doi: 10.4321/S1134-928X2014000200002.
- [2] Resnick B, Klinedinst NJ, Yerges-Armstrong L, Choi EY, Dorsey SG. The impact of genetics on physical resilience and successful aging. *J Aging Health*. 2015;27(6):1084–104; doi: 10.1177/0898264315577586.
- [3] Urtamo A, Jyväkorpi SK, Strandberg TE. Definitions of successful ageing: a brief review of a multidimensional concept. *Acta Biomed*. 2019;90(2):359–63; doi: 10.23750/abm.v90i2.8376.
- [4] Lowry KA, Vallejo AN, Studenski SA. Successful aging as a continuum of functional independence: lessons from physical disability models of aging. *Aging Dis*. 2012;3(1):5–15.
- [5] Krawczyk-Suszek M, Kleinrok A. Health-related quality of life (HRQoL) of people over 65 years of age. *Int J Environ Res Public Health*. 2022;19(2):625; doi: 10.3390/ijerph19020625.
- [6] Enssle F, Kabisch N. Urban green spaces for the social interaction, health and well-being of older people – an integrated view of urban ecosystem services and socio-environmental justice. *Environ Sci Policy*. 2020;109:36–44; doi: 10.1016/j.envsci.2020.04.008.
- [7] Lin Y-H, Chen Y-C, Tseng Y-C, Tsai S-T, Tseng Y-H. Physical activity and successful aging among middle-aged and older adults: a systematic review and meta-analysis of cohort studies. *Aging*. 2020;12(9):7704–16; doi: 10.18632/aging.103057.
- [8] Parra-Rizo MA, Vásquez-Gómez J, Álvarez C, Díaz-Martínez X, Troncoso C, Leiva-Ordoñez AM, Zapata-Lamana R, Cigarroa I. Predictors of the level of physical activity in physically active older people. *Behav Sci*. 2022;12(9):331; doi: 10.3390/bs12090331.
- [9] WHO. Ageing and health. Available from: <https://www.who.int/es/news-room/fact-sheets/detail/ageing-and-health> (accessed: 22.08.2022).
- [10] Booth FW, Roberts CK, Laye MJ. Lack of exercise is a major cause of chronic diseases. *Compr Physiol*. 2012;2(2):1143–211; doi: 10.1002/cphy.c110025.
- [11] Novotová K, Pavlů D, Dvořáčková D, Arnal-Gómez A, Espí-López GV. Influence of walking as physiological training to improve respiratory parameters in the elderly population. *Int J Environ Res Public Health*. 2022;19(13):7995; doi: 10.3390/ijerph19137995.
- [12] Vidal-Domínguez M, Labeaga-Azcona J, Casado-Durandez P, Madrigal-Muñoz A, López-Doblas J, Montero-Navarro A, Meli-Landwerlin G. Report 2016: The elderly in Spain. Statistical data by state and autonomous communities [in Spanish]. Available from: <https://dialnet.unirioja.es/servlet/libro?codigo=732091> (accessed: 21.11.2023).
- [13] WHO. Physical activity. Available from: <https://www.who.int/news-room/fact-sheets/detail/physical-activity> (accessed: 21.11.2023).
- [14] Cruz-Jentoft AJ, Baeyens JP, Bauer JM, Boirie Y, Cederholm T, Landi F, Martin FC, Michel J-P, Rolland Y, Schneider SM, Topinková E, Vandewoude M, Zamboni M; European Working Group on Sarcopenia in Older People. Sarcopenia: European consensus on definition and diagnosis: Report of the European working group on sarcopenia in older people. *Age Ageing*. 2010;39(4):412–23; doi: 10.1093/ageing/afq034.
- [15] Sáez-López M, Sánchez-Hernández N, Jiménez-Mola S, Alonso-García N, Valverde-García J. Assessment of pain in the elderly [in Spanish]. *Rev. Soc. Esp. Dolor*. 2015;22(6):271–4; doi: 10.4321/S1134-80462015000600008.
- [16] Welsh TP, Yang AE, Makris UE. Musculoskeletal pain in older adults: a clinical review. *Med Clin North Am*. 2020;104(5):855–72; doi: 10.1016/j.mcna.2020.05.002.
- [17] Burtscher J, Millet GP, Gatterer H, Vonbank K, Burtscher M. Does regular physical activity mitigate the age-associated decline in pulmonary function?. *Sports Med*. 2022;52(5):963–70; doi: 10.1007/s40279-022-01652-9.
- [18] Groessl EJ, Kaplan RM, Rejeski WJ, Katula JA, Glynn NW, King AC, Anton SD, Walkup M, Lu C-J, Reid K, Spring B, Pahor m. Physical activity and performance impact long-term quality of life in older adults at risk for major mobility disability. *Am J Prev Med*. 2019;56(1):141–6; doi: 10.1016/j.amepre.2018.09.006.
- [19] World Medical Association. World Medical Association Declaration of Helsinki: Ethical Principles for Medical Research Involving Human Subjects. *JAMA*. 2013;310(20):2191–4; doi: 10.1001/jama.2013.281053.
- [20] von Elm E, Altman DG, Egger M, Pocock SJ, Gøtzsche PC, Vandenbroucke JP; STROBE Initiative. The Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) statement: guidelines for reporting observational studies. *J Clin Epidemiol*. 2008;61(4):344–9; doi: 10.1016/j.jclinepi.2007.11.008.
- [21] United Nations. Ageing, older persons and the 2030 agenda for sustainable development. Available from: https://www.un.org/development/desa/dspd/wp-content/uploads/sites/22/2017/08/Ageing-Older-Persons-and-2030-Agenda_Issues-Brief-low-resolution-.pdf (accessed: 13.03.2025).


- [22] Gopinath B, Kifley A, Flood VM, Mitchell P. Physical activity as a determinant of successful aging over ten years. *Sci Rep.* 2018;8(1):10522; doi: 10.1038/s41598-018-28526-3.
- [23] Mantilla Toloza SC, Gómez-Conesa A. International Physical Activity Questionnaire. An adequate instrument in population physical activity monitoring. *Rev Iberoam Fisioter Kinesiol.* 2007;10(1):48–52; doi: 10.1016/S1138-6045(07)73665-1.
- [24] Geeta A, Jamaiyah H, Safiza MN, Khor GL, Kee CC, Ahmad AZ, Suzana S, Rahmah R, Faudzi A. Reliability, technical error of measurements and validity of instruments for nutritional status assessment of adults in Malaysia. *Singapore Med J.* 2009;50(10):1013–8.
- [25] Shah CH, Brown JD. Reliability and validity of the Short-Form 12 item version 2 (SF-12v2) health-related quality of life survey and disutilities associated with relevant conditions in the U.S. older adult population. *J Clin Med.* 2020;9(3):661; doi: 10.3390/jcm9030661.
- [26] Enríquez-Reyna MC, Cruz-Castruita RM, Ceballos-Gurrola O, García-Cadena CH, Hernández-Cortés PL, Guevara-Valtier MC. Psychometric properties of the Exercise Benefits/Barriers Scale in Mexican elderly women. *Rev Lat Am Enfermagem.* 2017;25:e2902; doi: 10.1590/1518-8345.1566.2902.
- [27] Hawker GA, Mian S, Kendzerska T, French M. Measures of adult pain: Visual Analog Scale for Pain (VAS Pain), Numeric Rating Scale for Pain (NRS Pain), McGill Pain Questionnaire (MPQ), Short-Form McGill Pain Questionnaire (SF-MPQ), Chronic Pain Grade Scale (CPGS), Short Form-36 Bodily Pain Scale (SF-36 BPS), and Measure of Intermittent and Constant Osteoarthritis Pain (ICOAP). *Arthritis Care Res.* 2011;63(Suppl 11):240–52; doi: 10.1002/acr.20543.
- [28] Bijur PE, Silver W, Gallagher EJ. Reliability of the visual analog scale for measurement of acute pain. *Acad Emerg Med.* 2001;8(12):1153–7; doi: 10.1111/j.1553-2712.2001.tb01132.x.
- [29] Springer BA, Marin R, Cyhan T, Roberts H, Gill NW. Normative values for the unipedal stance test with eyes open and closed. *J Geriatr Phys Ther.* 2007;30(1):8–15; doi: 10.1519/00139143-200704000-00003.
- [30] Hertel J, Miller SJ, Denegar CR. Intratester and intertester reliability during the star excursion balance tests. *J Sport Rehabil.* 2000;9(2):104–16; doi: 10.1123/jsr.9.2.104.
- [31] Powden CJ, Dodds TK, Gabriel EH. The reliability of the Star Excursion Balance test and Lower Quarter Y-Balance test in healthy adults: a systematic review. *Int J Sports Phys Ther.* 2019;14(5):683–94.
- [32] Giménez M, Servera E, Vergara P. Prevention and Rehabilitation in Chronic Respiratory Disease. *Physiotherapy, Training and Respiratory Care* [in Spanish]. Editorial Médica Panamericana; 2001.
- [33] Fernández-Villar A, Represas-Represas C, Mouronte-Roibás C, Ramos-Hernández C, Priegue-Carrera A, Fernández-García S, López-Campos JL. Reliability and usefulness of spirometry performed during admission for COPD exacerbation. *PLOS ONE.* 2018;13(3):e0194983; doi: 10.1371/journal.pone.0194983.
- [34] Holland AE, Spruit MA, Troosters T, Puhan MA, Pepin V, Saey D, McCormack MC, Carlin BW, Sciruba FC, Pitta F, Wanger J, MacIntyre N, Kaminsky DA, Culver BH, Revill SM, Hernandez NA, Andrianopoulos V, Camillo CA, Mitchell KE, Lee AL, Hill CJ, Singh SJ. An official European Respiratory Society/American Thoracic Society technical standard: field walking tests in chronic respiratory disease. *Eur Respir J.* 2014;44(6):1428–46; doi: 10.1183/09031936.00150314.
- [35] Pereiro AX, Campos-Magdaleno M, Navarro-Pardo E, Juncos-Rabadán O, Facal D. Normative scores for the Timed Up & Go in a Spanish sample of community-dwelling adults with preserved functionality. *Aten Primaria.* 2021;53(7):102065; doi: 10.1016/j.aprim.2021.102065.
- [36] Zhou J, Yao Q, Han R, De Bock P, Vassard-Yu G, Halle-mans A, Van Laer L. Reliability and validity of instrumented Timed Up and Go test in typical adults and elderly: a systematic review. *Arch Phys Med Rehabil.* 2005;106(7):1092–107; doi: 10.1016/j.apmr.2025.03.001.
- [37] Wanderley FAC, Silva G, Marques E, Oliveira J, Mota J, Carvalho J. Associations between objectively assessed physical activity levels and fitness and self-reported health-related quality of life in community-dwelling older adults. *Qual Life Res.* 2011;20(9):1371–8; doi: 10.1007/s11136-011-9875-x.
- [38] Ramos-Cordero P, Pinto-Fontanillo JA. Health status of the elderly. Current situation [in Spanish]. *Av Odon-toestomatol.* 2015;31(3):107–16; doi: 10.4321/S0213-12852015000300001.
- [39] Thomas ET, Guppy M, Straus SE, Bell KJL, Glasziou P. Rate of normal lung function decline in ageing adults: a systematic review of prospective cohort studies. *BMJ Open.* 2019;9(6):e028150; doi: 10.1136/bmjopen-2018-028150.
- [40] Pellegrino R, Viegi G, Brusasco V, Crapo RO, Burgos F, Casaburi R, Coates A, van der Grinten CPM, Gustafsson P, Hankinson J, Jensen R, Johnson DC, MacIntyre N, McKay R, Miller MR, Navajas D, Pedersen OF, Wanger J. Interpretative strategies for lung function tests. *Eur Respir J.* 2005;26(5):948–68; doi: 10.1183/09031936.05.00035205.
- [41] Vaz Fragoso CA, Gill TM. Respiratory impairment and the aging lung: a novel paradigm for assessing pulmonary function. *J Gerontol A Biol Sci Med Sci.* 2012;67(3):264–75; doi: 10.1093/gerona/glr198.
- [42] Poon ET-C, Wongpipit W, Ho RS-T, Wong SH-S. Interval training versus moderate-intensity continuous training for cardiorespiratory fitness improvements in middle-aged and older adults: a systematic review and meta-analysis. *J Sports Sci.* 2021;39(17):1996–2005; doi: 10.1080/02640414.2021.1912453.
- [43] Castañer M, Saüch G, Prat Q, Camerino O, Anguera MT. Perceived improvements in motor balance in relation to physical activity in programs for the elderly [in Spanish]. *Cuad Psicol Deporte.* 2016;16(1):77–84.
- [44] Dagnino APA, Campos MM. Chronic pain in the elderly: mechanisms and perspectives. *Front Hum Neurosci.* 2022;16:736688; doi: 10.3389/fnhum.2022.736688.
- [45] WHO. Mental health of older adults. Available from: <https://www.who.int/news-room/fact-sheets/detail/mental-health-of-older-adults> (accessed 03.11.2022).
- [46] Coupland C, Dhiman P, Morriss R, Arthur A, Barton G, Hippisley-Cox J. Antidepressant use and risk of adverse outcomes in older people: population based cohort study. *BMJ.* 2011;343:d4551; doi: 10.1136/bmj.d4551.
- [47] Staples WH, Kays A, Richman R. Examination of the correlation between physical and psychological measures in community-dwelling older adults. *Clin Interv Aging.* 2020;15:293–300; doi: 10.2147/CIA.S239053.
- [48] Cheong CY, Yap P, Gwee X, Chua DQL, Wee SL, Yap KB, Ng TP. Physical and functional measures predicting long-term mortality in community-dwelling older adults: a comparative evaluation in the Singapore Longitudinal Ageing Study. *Aging.* 2021;13(23):25038–54; doi: 10.18632/aging.203756.

[49] Mulasso A, Roppolo M, Rabaglietti E. The role of individual characteristics and physical frailty on health related quality of life (HRQOL): a cross sectional study of Italian community-dwelling older adults. Arch Gerontol Geriatr. 2014;59(3):542–8; doi: 10.1016/j.archger.2014.08.012.

[50] Salguero A, Martínez-García R, Molinero O, Márquez S. Physical activity, quality of life and symptoms of depression in community-dwelling and institutionalized older adults. Arch Gerontol Geriatr. 2011;53(2):152–7; doi: 10.1016/j.archger.2010.10.005.

[51] Moreira B de S, Dos Anjos DM da C, Pereira DS, Dias RC, Kirkwood RN. The geriatric depression scale and the timed up and go test predict fear of falling in community-dwelling elderly women with type 2 diabetes mellitus: a cross-sectional study. BMC Geriatr. 2016;16:56; doi: 10.1186/s12877-016-0234-1.

Supplementary material 1 (A) percentage of participants with pain in different body areas
 (B) pain intensity of each of the painful body areas



Supplementary material 2

Correlations between physical activity, pain, balance, respiratory and functional outcomes, and quality of life

Variables	Physical health-related quality of life	Mental health-related quality of life	r	p
	r	p		
Physical activity (METs/week)	0.25*	0.02	0.27*	0.01
Musculoskeletal pain (VAS)	-0.43**	< 0.001	-0.24*	0.03
BALANCE				
UPST (s)				
right leg	0.26*	0.01	0.09	0.41
left leg	0.08	0.45	0.12	0.25
YBT-LQ (cm)				
right leg	0.16	0.14	0.11	0.34
left leg	0.19	0.09	0.07	0.55
Cardiorespiratory variables				
HR (bpm)	-0.02	0.85	-0.20	0.06
SpO ₂	0.29**	0.01	0.13	0.23
FVC (l)	0.37**	< 0.001	0.34**	0.001
FVC (%) predicted	0.33**	0.002	0.13	0.23
FEV ₁ (l/min)	0.36**	< 0.001	0.30**	0.01
FEV ₁ (%) predicted	0.31**	0.003	0.15	0.16
FEV ₁ /FVC (%)	0.01	0.96	0.07	0.55
FUNCTIONAL VARIABLES				
6MWT				
distance (m)	0.27*	0.01	0.27*	0.01
SpO ₂	0.23*	0.03	0.07	0.53
HR (bpm)	0.11	0.32	-0.07	0.52
dyspnoea	0.01	0.96	0.04	0.72
TUG (s)	-0.23*	0.03	-0.32**	0.002
Quality of life				
SF-12 physical subscale	-	-	0.36**	0.001
SF-12 mental subscale	0.36**	0.001	-	-

Data shown as Pearson correlation coefficient.

SF-12 – Short Form 12 Health Survey, MET – Metabolic Equivalent of Task, VAS – Visual Analogue Scale, HR – heart rate, SpO₂ – oxygen saturation, FVC – forced vital capacity, FEV₁ – forced expiratory volume in one second, UPST – Unipedal Stance Test, YBT-LQ – the Lower Quarter Y-Balance Test, 6MWT – Six-Minute Walk Test, TUG – Timed Up and Go

* p < 0.05, ** p < 0.01