

Study of the effectiveness of dry needling in the context of treatment of low back pain in a post-surgical patient: a case study

DOI: <https://doi.org/10.5114/pq/217016>

Aleksander Burzawa¹ , Mateusz Stefański² , Mattia Di Stefani¹ , Julia Kuberska¹ , Tomasz Król³ , Michał Kuszewski² 

¹ Student's research science team FASCYNaci, Academy of Physical Education in Katowice, Katowice, Poland

² Institute of Physiotherapy, Academy of Physical Education in Katowice, Katowice, Poland

³ Department of Kinesitherapy and Special Methods, School of Health Sciences in Katowice, Medical University of Silesia, Katowice, Poland

Abstract

Introduction. The purpose of the study is to investigate the effectiveness of dry needling (DN) in the context of treating LBP (low back pain)-related symptoms over a period of four weeks following a surgery of laparoscopic removal of an ovarian cyst in a female patient.

Methods. The Visual Analogue Scale (VAS) was chosen to assess the patient's pain level, and functional status was assessed using the Oswestry Disability Index (ODI). An inclinometer was applied to measure the ranges of motion in the lumbar spine in the sagittal and frontal planes, and functional capacity was evaluated using the Y-Balance test. Soft tissue properties in the area of the lumbar spine were checked for mobility in the sagittal plane using myotonometry. The muscles punctured as part of therapy included the erector spinae, gluteus maximus, gluteus medius, and piriformis.

Results. The patient's reported pain level decreased. Her functional status, assessed according to the ODI questionnaire, improved. Inclinometric examination did not reveal obvious changes for most of the ranges of motion tested. Assessment of the right lower limb functional capacity with regard to the frontal, posteromedial, and posterolateral planes using the Y-Balance test also showed no significant changes. As for the myotonometric test, the greatest differences were noted in the dynamic stiffness parameter (S), present over almost the entire targeted area. A similar correlation was found in the parameters of relaxation (R) and creep (C).

Conclusions. The use of a dry needling procedure promotes tone symmetrisation in soft tissues and reduces the level of pain experienced.

Key words: myotonometry, pain reduction, soft tissue, muscles

Introduction

Low back pain (LBP) refers to the pain felt between the lower border of the ribs and the intergluteal cleft [1]. LBP in itself is not a disease but a symptom whose origin is often difficult to determine [1].

Even though LBP problems affect a wide range of populations of all ages, from teenagers to the elderly, the condition clearly becomes more common with age [1–4]. It is believed that the majority of the adult population will experience LBP at a certain stage in their lives [2, 4]. LBP is also considered the most common cause of disability [1–3]. The condition is as common in highly developed countries as it is in developing or underdeveloped ones. Women suffer from the condition more frequently and thus are more likely to seek medical help to improve their health [1–4].

In most patients, it seems impossible to determine the specific cause of LBP [2–4]. Imaging studies such as magnetic resonance imaging (MRI) or computed tomography (CT) performed on patients with LBP often display various anatomical abnormalities, but the same defects are also often found in patients who do not report pain [1]. In young patients, the presence of certain abnormalities on imaging studies may increase the chance of LBP occurring in the future [1].

It is assumed that LBP may be caused by a combination of several factors, including biophysical features (muscle mass, composition or volume, anatomical structure), psychological factors (fear, depression) and social aspects (level of education, access to health care, quality of health care, occupation, financial situation, etc.) [2]. These complaints are sometimes caused by specific conditions, such as vertebral fractures, neoplastic lesions, or inflammatory diseases, but these require specific treatment and constitute a small percentage of LBP cases [2–4].

A method that is gaining popularity as a treatment option for LBP is dry needling (DN). The treatment involves identifying and then puncturing active myofascial trigger points (MTrPs) in skeletal muscles. This causes a local twitch response (LTR) in the tense muscle band, resulting in its relaxation [5]. A puncture is done with a thin acupuncture needle, but the method is strictly associated with conventional Western medicine [6]. DN is a minimally invasive method, and it is also simple to perform [5]. It can be effective in providing pain relief, and it improves the functional status of the patient [5, 7]. Despite the potential benefits of DN in the treatment of LBP, the number of research studies and publications on the therapy is low, and the available articles describing positive effects of the treatment emphasise the need of further research into the effectiveness of DN in LBP [7, 8].

Correspondence address: Tomasz Król, Department of Kinesitherapy and Special Methods, School of Health Sciences in Katowice, Medical University of Silesia, Medyków 12 street, 40-752 Katowice, Poland, e-mail: tkrol@sum.edu.pl; <https://orcid.org/0000-0002-0954-5117>

Received: 04.07.2025

Accepted: 15.01.2026

Citation: Burzawa A, Stefański M, Di Stefani M, Kuberska J, Król T, Kuszewski M. Study of the effectiveness of dry needling in the context of treatment of low back pain in a post-surgical patient: a case study. *Physiother Quart.* 2026;34(3):120–125; doi: <https://doi.org/10.5114/pq/217016>.

The purpose of the present study is to investigate the effectiveness of dry needling (DN) in the context of treating LBP-related symptoms over a period of four weeks following laparoscopic removal of an ovarian cyst in a female patient.

Subjects and methods

The patient is a female aged 24, with a self-reported height of 182 cm and a body weight of 72 kg (BMI 21.74 kg/m² – within normal limits). The average level of physical activity per week is, according to the patient’s declaration, 2 hours. The patient has no additional comorbidities or addictions. She underwent a laparoscopic surgery to remove an ovarian cyst 2 years ago. She reports back pain in the lumbar area, which persists at an average of 5 points on the Numeric Rating Scale (NRS) scale and started approximately 4 years ago. The patient was not taking any painkiller medication during the course of this study.

The inclusion criteria were the following: age 20 to 30 years, low back pain rated at 4 to 9 on the NRS persisting for at least six weeks, and a history of abdominal surgery a minimum of two years ago.

The exclusion criteria were a lack of consent to participate in the study and contradictions to the dry needling procedure.

The project was designed as an observer-blinded trial, where blinding was achieved by separating the study from the therapy. The researchers did not know whether or what kind of therapy the patient was undergoing; the person performing the therapy, in turn, did not know the results of the patient’s subsequent tests.

Measuring tools

The patient’s functional status was assessed using the Oswestry Disability Index (ODI). This questionnaire is divided into ten categories, including pain severity, pain while lifting, sitting, sleeping, travelling, daily care, walking, standing, socialising, as well as the assessment of changes in pain severity. Each category is scored from 0 to 5 points, giving a maximum of 50 points. The score ranges for particular results are established as follows: 0 to 4 points – no disability, 5 to 14 points – mild disability, 15 to 24 points – moderate disability, 25 to 34 points – severe disability, 35 to 50 points – total disability [9].

In order to assess the patient’s pain level, a NRS of 0 to 10 was selected, where 0 means the absence of pain, and 10 means the highest imaginable pain level [10].

The ranges of motion of the lumbar spine in the sagittal and frontal planes were measured using a Saunders digital inclinometer (Baseline Digital Inclinometer, The Saunders Group Inc, Chaska, MN, USA) [11]. The Y-Balance test was used to assess functional capacities in the anterior, posterolateral, and posteromedial planes for the patient’s lower body [12].

In addition, the viscoelastic properties of soft tissues in the lumbar and hip sections were assessed. The points at which the measurements were made were the so-called coordination centres and were taken from the concept of Fascial Manipulation®. These were the points: AN and RE Lu and Cx (both sides) anatomically. It was 8 points in total (4 for the Lu segment and 4 for the Cx segment). These points, which lie on the rectus abdominis muscle (AN Lu) (lateral to the navel) and on the erector spinae muscle (RE Lu) (at the level of L1) and below the inguinal ligament (above the iliolumbar muscle) and on the gluteus maximus muscle (above the

ischial tuberosity, were evaluated for movements in the sagittal plane using myotonometry (MyotonPRO Tallinn, Estonia). The soft tissue properties measured included frequency (F), dynamic stiffness (S), decrement (D), time required for regaining tissue relaxation (R) and creep (C) [13–15]. The parameters were also evaluated in the area of abdominal scar tissue resulting from the surgical procedure.

Experiment protocol

Measurement procedures were carried out in the following sequence: assessment of pain levels using the NRS, assessment of functional status using the ODI questionnaire, evaluation of mobility ranges in the sagittal and frontal planes using an inclinometer, assessment of functional capacity in the anterior, posterolateral and posteromedial planes for the lower body using the Y-Balance test, and finally myotonometry.


As for inclinometric measurement, the result is the arithmetic average of three subsequent measurements of one movement in a given plane. In the case of the Y-Balance test, prior to the measurement, the patient performed six ‘warm-up’ repetitions for each plane, for each lower limb tested, in order to get familiar with the test being performed. Measurements were performed using an original Y-Balance test kit, and the patient was not wearing shoes at the time of testing. The result of the Y-Balance test is the arithmetic average of three measurements taken in each plane for each lower extremity.

Measurement procedures were carried out immediately before the first therapy session, and the post-test was conducted immediately after the last therapy session. Both measurements were performed by the same person at similar times of the day and under similar conditions.

Figure 1 presents the experiment as a flowchart.

Therapeutic procedures

DN therapy was performed using needles by yellowSPORT (Katowice, Poland) made of 0.3-mm-thick surgical steel. The



length of the needle to be used was selected based on the puncture site.

The muscles punctured as part of the therapy included the erector spinae, gluteus maximus, gluteus medius, and piriformis [7–8].

Prior to the application of the needles, the therapist disinfected their hands, put on disposable gloves and disinfected the puncture site. The therapist was unaware of the results of the patient’s functional test (observer-blinded trial).

The testing protocol was the following:

1. Erector spinae – after palpation of the spinous processes of the lumbar vertebrae, the insertions were performed along the fibres of the spinal erectors at the level of spinal segments L1 to L5; each needle was positioned caudomedially with an inclination of approximately 30°. The technique of stimulation was rhythmic ‘pistoning’ of the needle. After the insertion of the needles, the application was left undisturbed for 15 min, and then the muscle was stimulated again by ‘pistoning’ until the insertion was completed [7, 8]. The needle used was 25 mm long.

2. Gluteus medius and gluteus maximus muscles – after palpation of the iliac crest and tight bands within the bulk of the muscles, the insertion occurred perpendicularly to the skin surface in the region of increased tension. A stimulation technique of rhythmic ‘pistoning’ of the needle was utilised until LTR was achieved. After the needle’s insertion, the application was left undisturbed for 15 minutes, and then the muscle was again stimulated by ‘pistoning’ the needle until the insertion was completed [7, 8]. The needle used was 75 mm long.

3. Piriformis muscles – after palpation of the lateral border of the sacrum, the greater trochanter of the femur and tight bands within the bulk of the muscles, the insertion was applied perpendicularly to the skin surface in the region of increased tension. A stimulation technique of rhythmic ‘pistoning’ of the needle was utilised until LTR was achieved. After the needle’s insertion, the application was left undisturbed for 15 min, and then the muscle was again stimulated by ‘pistoning’ until the insertion was completed [7, 8]. The needle used was 75 mm long.

The therapy was carried out twice a week over a period of four weeks, with a minimum of two days’ breaks between therapy sessions in any given week [7, 8].

Results

The patient’s reported pain level decreased after the intervention; prior to the treatment, the patient rated the pain severity at 5 points on the NRS, whereas after the therapy,

the reported pain level was 2 points on the NRS. Figure 2 graphically demonstrates the result.

The patient’s functional status, assessed according to the ODI questionnaire, improved. Before the intervention, it was rated by the patient at 7 points. After the therapy, the score according to the ODI questionnaire was 3. Figure 3 graphically demonstrates the result, along with particular categories of the ODI questionnaire.




Figure 2. Pain level on the Numeric Rating Scale (NRS)




Figure 4. Range of motion in the spine – averaged results




Figure 3. Functional status according to the Oswestry Disability Index questionnaire

Table 1. Y-Balance test results for the right lower limb

Trial No.	Frontal plane		Posteromedial plane		Posterolateral plane	
	before therapy	after therapy	before therapy	after therapy	before therapy	after therapy
Trial 1	66	68	76	76	76	78
Trial 2	69	70	76	82	72	81
Trial 3	71	74	80	76	84	83
Averaged result	68.7	70.7	77.3	78	77.3	80.7

Table 2. Y-Balance test results for the left lower limb

Trial No.	Frontal plane		Posteromedial plane		Posterolateral plane	
	before therapy	after therapy	before therapy	after therapy	before therapy	after therapy
Trial 1	67	70	80	74	62	83
Trial 2	69	74	80	79	70	87
Trial 3	72	73	82	80	78	90
Averaged result	69.3	72.3	80.7	77.7	70	86.7

Table 3. Results of myotonometry – lumbar spine movements in the sagittal plane

Segment and kind of movement	F (Hz)		S (N/m)		D		R (ms)		C	
	before therapy	after therapy	before therapy	after therapy	before therapy	after therapy	before therapy	after therapy	before therapy	after therapy
Forward movement of the L. spine – right side	11.1	11.4	159	141	0.79	0.83	29.5	27.5	1.66	1.43
Forward movement of the L. spine – left side	12.0	12.2	132	164	1.00	0.90	26.6	26.5	1.30	1.48
Forward movement of the hip – right side	9.1	9.1	164	124	1.67	1.94	37.5	34.3	2.60	1.73
Forward movement of the hip – left side	9.3	9.3	131	123	1.47	1.72	36.9	34.4	2.00	1.74
Backward movement of the L. spine – right side	16.2	14.5	313	255	1.13	1.09	16.4	18.7	0.99	1.10
Backward movement of the L. spine – left side	14.7	15.4	227	256	1.03	0.97	18.2	18.7	1.00	1.10
Backward movement of the hip – right side	11.8	15.1	158	213	1.34	1.52	26.3	23.1	1.42	1.34
Backward movement of the hip – left side	10.8	15.2	144	205	1.30	1.47	31.1	23.6	1.69	1.38

F – frequency, S – stiffness, D – decrement, R – relaxation, C – creep

Table 4. Results of myotonometry in the area of postsurgical scar tissue

Place of measurement	F (Hz)		S (N/m)		D		R (ms)		C	
	before therapy	after therapy	before therapy	after therapy	before therapy	after therapy	before therapy	after therapy	before therapy	after therapy
Above the scar – right side	11.1	12.2	161	184	1.13	1.17	28.9	26.1	1.61	1.51
Right side of the scar – right side	11.3	12.1	158	188	1.01	1.18	28.5	26.8	1.55	1.60
Below the scar – right side	12.2	12.5	186	197	0.99	1.20	25.4	24.8	1.46	1.46
Left side of the scar – right side	12.9	13.3	195	212	1.15	1.22	22.7	22.6	1.27	1.31
Above the scar – left side	11.0	12.1	148	183	1.37	1.28	26.9	26.6	1.37	1.52
Right side of the scar – left side	12.9	13.1	212	205	1.23	1.20	21.6	21.9	1.24	1.24
Below the scar – left side	11.7	12.3	187	198	1.21	1.23	24.8	24.4	1.40	1.44
Left side of the scar – left side	11.3	12.1	179	178	1.23	1.36	33.0	27.5	1.79	1.56

F – frequency, S – stiffness, D – decrement, R – relaxation, C – creep

In the inclinometric test, there were no apparent changes for flexion, extension, and lateral flexion to the right. A significant change in the result was noted for lateral flexion to the left; the improvement in the averaged result exceeded 6 degrees. Figure 4 shows the averaged results of all the measurements. The numbers are presented in degrees.

Assessment of the right lower limb functional capacity for the frontal, posteromedial, and posterolateral planes with the use of the Y-Balance test also showed no obvious changes before and after the intervention. A significant change in the result was only noted for the posterolateral plane of the left lower limb: the improvement in the averaged result exceeded 16 cm. The results of particular measurements for the right and left lower extremities are presented in Tables 1 and 2, respectively. The results are presented in centimetres.

In the myotonometric test, the biggest differences are noted in the dynamic stiffness parameter (S) over almost the entire targeted area. A similar dependence can be observed in the parameters of relaxation (R) and creep (C). It is also worth noting the increased tonic frequency in the tissue (F) that results from the intervention. The results of the myotonometric study are presented in Tables 3 and 4.

Discussion

The results obtained in the present study suggest that therapeutic intervention of dry needling carried out in the course of LBP, in the form described in the article, can be an effective method of relieving low back pain. The results, as based on the patient's perceived pain experience (NRS and ODI questionnaire), indicate a significant improvement in this regard. Within the framework of the proposed methodology, no improvement was observed in the patient's functional capacity. The results obtained from myotonometric measurements suggest that the intervention through dry needling leads to soft tissues being toned and symmetrised.

The results indicative of pain reduction and tension symmetrisation within soft tissues are an excellent starting point for functional improvement in the patient. Perhaps such an effect might also have been achieved, but the very short time between the end of the therapeutic intervention and the repeat examination (measurement taken immediately after the end of the treatment) did not allow us to demonstrate such a correlation.

A study by Loizidis *et al.* [16] showed similar effects of dry needling interventions in patients with LBP. Patient-reported pain levels decreased markedly, and functional improvement was also noted in the patients undergoing the intervention.

DN therapy is also effective when combined with other physiotherapy treatments. Pain reduction in sciatic hyperalgesia has been observed after DN therapy combined with neurodynamic treatments [17].

As noted in the introduction, LBP refers to a wide range of conditions, and pain relief achieved through dry needling alone may not be sufficient in the overall treatment process. Medical functional training seems to be an appropriate complement to the proposed therapeutic intervention [18] as it clearly improves the patient's functional parameters. The use of dry needling as a means of reducing pain and normalising tissue properties could constitute a good starting point for the consolidation and progression of the achieved effects by implementing medical training.

Such conclusions have been drawn from research conducted by Martín-Corrales *et al.* [19]. According to this study, complementing a medical training program with dry needling may result in significantly better patient pain management

compared to a control group where dry needling was not introduced as part of the therapy.

Similar results regarding the reduction of pain in LBP patients were observed in a study conducted by Pawlukiewicz *et al.* [20]. In this study, a group of patients undergoing a standard therapy protocol based on fascial manipulation achieved a significant reduction in pain levels by an average of 3.5 points on the VAS scale after the intervention.

Myotonometric measurements conducted by Kim and Park [21] on the tissues closely associated with LBP (quadratus lumborum, gluteus medius, hamstring muscles) revealed a significant decrease in the dynamic stiffness of the tissues both in the group subjected to core stability training focused on the lumbar region and in the group who underwent resistance-based training and passive hip joint stretching. However, no significant differences were observed between the two groups. Thus, it seems that the inclusion of medical training in the therapeutic program may also help to normalise soft tissue tension.

The improvement in the patient's range of motion of spinal lateral flexion to the left appears to be reflected in the myotonometric measurements. The measurements taken in the lumbar area on the right side of the body showed that the application of dry needling resulted in the reduction of dynamic stiffness (S), time required to relax again (R) and creep (C). Perhaps the normalisation of tissue tension in this region allowed for a greater range of motion in this direction; however, it should be emphasised that this is the only tested plane of motion in which an improvement in the result was noted. A similar result was obtained in a study conducted by Lee and Lee [22]. In patients undergoing the procedure of medical functional training accompanied by manual therapy (Maitland's concept), the tone of the descending part of the trapezius muscle was significantly decreased, and the range of motion of the cervical spine improved significantly in all directions.

Limitations

Importantly, the present study does not include catamnesis; hence it is impossible to determine how long the effect obtained by applying the intervention would continue. Moreover, since the study was conducted on only one patient, it is infeasible to determine whether the outcome obtained by applying the proposed intervention would be equally effective for a broader population. Nevertheless, this may be expected, given the extent of changes observed in this single case and the analysis of literature reports.

Conclusions

Based on the analysis of the results of the study, the following conclusions have been drawn:

1. The use of the dry needling procedure has an impact on the patient's pain level – a significant decrease is observed.
2. The use of the dry needling procedure has no immediate effect on improving the patient's functional status, yet it can be a good starting point for achieving this effect should the therapeutic process be complemented with, for instance, medical functional training.
3. The use of the dry needling procedure has an impact on the tension symmetry of the soft tissues.

Ethical approval

The research related to human use complied with all the relevant national regulations and institutional policies, followed the tenets of the Declaration of Helsinki, and was ap-

proved by the Ethics Committee of the Physiotherapy Poland Association (approval No.: 2/07/2024).

Informed consent

Informed consent has been obtained from all individuals included in this study.

Disclosure statement

No author has any financial interest or received any financial benefit from this research.

Conflict of interest

The authors state no conflict of interest.

Funding

This research received no external funding.

References

- [1] Buchbinder R, van Tulder M, Öberg B, Costa LM, Woolf A, Schoene M, Croft P; Lancet Low Back Pain Series Working Group. Low back pain: a call for action. *Lancet*. 2018;391(10137):2384–8; doi: 10.1016/S0140-6736(18)30488-4.
- [2] Hartvigsen J, Hancock MJ, Kongsted A, Louw Q, Ferreira ML, Genevay S, Hoy D, Karppinen J, Pransky G, Sieper J, Smeets RJ, Underwood M; Lancet Low Back Pain Series Working Group. What low back pain is and why we need to pay attention. *Lancet*. 2018;391(10137):2356–67; doi: 10.1016/S0140-6736(18)30480-X.
- [3] Maher C, Underwood M, Buchbinder R. Non-specific low back pain. *Lancet*. 2017;389(10070):736–47; doi: 10.1016/S0140-6736(16)30970-9.
- [4] Urits I, Burshtein A, Sharma M, Testa L, Gold PA, Orhuru V, Viswanath O, Jones MR, Sidransky MA, Spektor B, Kaye AD. Low back pain, a comprehensive review: pathophysiology, diagnosis, and treatment. *Curr Pain Headache Rep*. 2019;23:1-10.
- [5] Liu L, Huang Q-M, Liu Q-G, Thitham N, Li L-H, Ma Y-T, Zhao J-M. Evidence for dry needling in the management of myofascial trigger points associated with low back pain: a systematic review and meta-analysis. *Arch Phys Med Rehabil*. 2018;99(1):144–52.e2; doi: 10.1016/j.apmr.2017.06.008.
- [6] Kalichman L, Vulfsons S. Dry needling in the management of musculoskeletal pain. *J Am Board Fam Med*. 2010;23(5):640–6; doi: 10.3122/jabfm.2010.05.090296.
- [7] Rajfur J, Rajfur K, Kosowski Ł, Walewicz K, Dymarek R, Ptazkowski K, Taradaj J. The effectiveness of dry needling in patients with chronic low back pain: a prospective, randomized, single-blinded study. *Sci Rep*. 2022; 12(1):15803; doi: 10.1038/s41598-022-19980-1.
- [8] Rajfur J, Rajfur K, Matusz T, Malarska M, Walewicz K, Ptazkowski K, Dymarek R, Taradaj J. Dry needling with the use of FRSc technique in addition to standard rehabilitation program for chronic low back pain: a randomized controlled trial using both PROMs and measurement tools. *J Pain Res*. 2024;17:1041–53; doi: 10.2147/JPR.S450119.
- [9] Chiarotto A, Maxwell LJ, Terwee CB, Wells GA, Tugwell P, Ostelo RW. Roland-Morris Disability Questionnaire and Oswestry Disability Index: which has better measurement properties for measuring physical functioning in nonspecific low back pain? Systematic review and meta-analysis. *Phys Ther*. 2016;96(10):1620–37; doi: 10.2522/ptj.20150420.
- [10] Haefeli M, Elfering A. Pain assessment. *Eur Spine J*. 2006; 15(Suppl 1):17–24; doi: 10.1007/s00586-005-1044-x.
- [11] Saur PM, Ensink FBM, Frese K, Seeger D, Hildebrandt J. Lumbar range of motion: reliability and validity of the inclinometer technique in the clinical measurement of trunk flexibility. *Spine*. 1996;21(11):1332–8; doi: 10.1097/00007632-199606010-00011.
- [12] Alshehri Y, Alkhathami K, Brizzolaro K, Weber M, Wang-Price S. Reliability and validity of the Y-balance test in young adults with chronic low back pain. *Int J Sports Phys Ther*. 2021;16(3):628–35; doi: 10.26603/001c.23430.
- [13] Vatovec R, Voglar M. Changes of trunk muscle stiffness in individuals with low back pain: a systematic review with meta-analysis. *BMC Musculoskelet Disord*. 2024;25(1): 155; doi: 10.1186/s12891-024-07241-3.
- [14] Li Y, Yu J, Zhang J, Wang X. Quantifying the stiffness of lumbar erector spinae during different positions among participants with chronic low back pain. *PLOS ONE*. 2022;17(6):e0270286; doi: 10.1371/journal.pone.0270286.
- [15] Lo WL, Yu Q, Mao Y, Li W, Hu C, Li L. Lumbar muscles biomechanical characteristics in young people with chronic spinal pain. *BMC Musculoskelet Disord*. 2019;20: 559; doi: 10.1186/s12891-019-2935-z.
- [16] Loizidis T, Nikodelis T, Bakas E, Kollias I. The effects of dry needling on pain relief and functional balance in patients with sub-chronic low back pain. *J Back Musculoskelet Rehabil*. 2020;33(6):953–9; doi: 10.3233/BMR-181265.
- [17] Shanmugam S, Kandakurti PK, Durairaj S. The effectiveness of deep dry needling combined with neural mobilisation in patients with sciatic hyperalgesia: a case report. *Physiother Quart*. 2025;33(3):42–6; doi: 10.5114/pq/191031.
- [18] Cortell-Tormo JM, Sánchez PT, Chulvi-Medrano I, Tortosa-Martínez J, Manchado-López C, Llana-Belloch S, Pérez-Soriano P. Effects of functional resistance training on fitness and quality of life in females with chronic non-specific low-back pain. *J Back Musculoskelet Rehabil*. 2018;31(1):95–105; doi: 10.3233/BMR-169684.
- [19] Martín-Corrales C, Bautista IV, Méndez-Mera JE, Fernández-Matías R, Achalandabaso-Ochoa A, Gallego-Izquierdo T, Nuñez-Nagy S, Pecos-Martín D. Benefits of adding gluteal dry needling to a four-week physical exercise program in a chronic low back pain population. A randomized clinical trial. *Pain Med*. 2020;21(11):2948–57; doi: 10.1093/pm/pnaa279.
- [20] Pawlukiewicz M, Kochan M, Niewiadomy P, Szuścik-Niewiadomy K, Taradaj J, Król P, Kuszewski MT. Fascial manipulation method is effective in the treatment of myofascial pain, but the treatment protocol matters: a randomized control trial – preliminary report. *J Clin Med*. 2022;11(15):4546; doi: 10.3390/jcm11154546.
- [21] Kim J-H, Park H-Y. Effects of combined hip exercise and passive stretching on muscle stiffness, pain perception and pain-related disability, and physical function in older adults with low back pain. *Phys Act Nutr*. 2022;26(3): 16–24; doi: 10.20463/pan.2022.0014.
- [22] Lee K-S, Lee J-H. Effect of Maitland mobilization in cervical and thoracic spine and therapeutic exercise on functional impairment in individuals with chronic neck pain. *J Phys Ther Sci*. 2017;29(3):531–5; doi: 10.1589/jpts.29.531.

Copyright: © 2026 Wrocław University of Health and Sport Sciences. This is an Open Access journal distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs (CC BY-NC-ND) (<https://creativecommons.org/licenses/by-nc-nd/4.0/legalcode>), allowing third parties to download and share its works but not commercially purposes or to create derivative works.